

Global well-posedness and scattering for the mass-critical Hartree equation with radial data

Changxing Miao[†] Guixiang Xu[†] and Lifeng Zhao[‡]

[†]Institute of Applied Physics and Computational Mathematics
P. O. Box 8009, Beijing, China, 100088

[‡] Department of Mathematics, University of Science and Technology of China
(miao_changxing@iapcm.ac.cn, xu_guixiang@iapcm.ac.cn, zhao_lifeng@iapcm.ac.cn)

Abstract: We establish global well-posedness and scattering for solutions to the mass-critical nonlinear Hartree equation $iu_t + \Delta u = \pm(|x|^{-2} * |u|^2)u$ for large spherically symmetric $L_x^2(\mathbb{R}^d)$ initial data; in the focusing case we require, of course, that the mass is strictly less than that of the ground state.

Résumé: Nous établissons l'existence globale et la diffusion des solutions de l'équation non linéaire de masse critique de Hartree $iu_t + \Delta u = \pm(|x|^{-2} * |u|^2)u$ pour des données initiales grandes à symétrie sphérique dans $L_x^2(\mathbb{R}^d)$; dans le cas focalisant nous imposons, bien sûr, que la masse soit strictement inférieure à celle de l'état fondamental.

Key Words: Hartree equation, global well-posedness, scattering, mass-critical.

AMS Classification: 35Q40, 35Q55, 47J35.

1 Introduction

We primarily consider the mass-critical Hartree equation:

$$\begin{cases} iu_t + \Delta u = \mu(|x|^{-2} * |u|^2)u, & \text{in } \mathbb{R}^d \times \mathbb{R}, \\ u(0) = u_0(x), & \text{in } \mathbb{R}^d. \end{cases} \quad (1)$$

where $d \geq 3$, $\mu = \pm 1$, with $\mu = +1$ known as the defocusing case and $\mu = -1$ as the focusing case. The Hartree equation arises in the study of Boson stars and other physical phenomena, see for example [18]. In chemistry, it appears as a continuous-limit model for mesoscopic molecular structures, see [7].

Definition 1.1. A function $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ on a non-empty time interval $I \subset \mathbb{R}$ (possibly infinite or semi-infinite) is a strong $L^2(\mathbb{R}^d)$ solution to (1) if it lies in the class

$C_t^0 L_x^2(K \times \mathbb{R}^d) \cap L_t^6 L_x^{\frac{6d}{3d-2}}(K \times \mathbb{R}^d)$ for all compact $K \subset I$, and we have the Duhamel formula

$$u(t_1) = e^{i(t_1-t_0)\Delta} u(t_0) - i \int_{t_0}^{t_1} e^{i(t_1-t)\Delta} F(u(t)) dt \quad (2)$$

for all $t_0, t_1 \in I$, where $F(u) = \mu(|x|^{-2} * |u|^2)u$. We refer to the interval I as the lifespan of u . We say that u is a maximal-lifespan solution if the solution cannot be extended to any strictly larger interval. We say that u is a global solution if $I = \mathbb{R}$.

Definition 1.2. We say that a solution u to (1) blows up forward in time if there exists a time $t_0 \in I$ such that

$$\|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}([t_0, \sup(I)) \times \mathbb{R}^d)} = \infty \quad (3)$$

and that u blows up backward in time if there exists a time $t_0 \in I$ such that

$$\|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}((\inf(I), t_0] \times \mathbb{R}^d)} = \infty. \quad (4)$$

For the mass-critical Hartree equation (1) with data in L^2 , the authors obtained some well-posedness and scattering results in [13] using the method of [5]. We collect these facts as follows:

Theorem 1.1 (Local well-posedness). *Given $u_0 \in L_x^2(\mathbb{R}^d)$ and $t_0 \in \mathbb{R}$, there exists a unique maximal-lifespan solution u to (1) with $u(t_0) = u_0$. We will write I for the maximal lifespan. This solution also has the following properties:*

- ★ (Local existence) I is an open neighborhood of t_0 .
- ★ (Mass conservation) The solution u has a conserved mass: for $t \in I$,

$$M(u) = M(u(t)) := \int_{\mathbb{R}^d} |u(t, x)|^2 dx. \quad (5)$$

- ★ (Blowup criterion) If $\sup(I)$ is finite, then u blows up forward in time; if $\inf(I)$ is finite, then u blows up backward in time.

- ★ (Continuous dependence) If $u_0^{(n)}$ is a sequence converging to u_0 in $L_x^2(\mathbb{R}^d)$ and $u^{(n)} : I^{(n)} \times \mathbb{R}^d \rightarrow \mathbb{C}$ are the associated maximal-lifespan solutions, then $u^{(n)}$ converges locally uniformly to u .

- ★ (Scattering) If u does not blow up forward in time, then $\sup(I) = +\infty$ and u scatters forward in time, that is, there exists a unique $u_+ \in L_x^2(\mathbb{R}^d)$ such that

$$\lim_{t \rightarrow +\infty} \|u(t) - e^{it\Delta} u_+\|_{L_x^2(\mathbb{R}^d)} = 0.$$

Similarly, if u does not blow up backward in time, then $\inf(I) = -\infty$ and u scatters backward in time, that is, there is a unique $u_- \in L_x^2(\mathbb{R}^d)$ so that

$$\lim_{t \rightarrow -\infty} \|u(t) - e^{it\Delta} u_-\|_{L_x^2(\mathbb{R}^d)} = 0.$$

- ★ (Spherical symmetry) If u_0 is spherically symmetric, then u remains spherically symmetric for all time.

★ *(Small data global existence)* If $M(u_0)$ is sufficiently small, then u is a global solution which does not blow up either forward or backward in time. Indeed, in this case

$$\int_{\mathbb{R}} \left(\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \lesssim M(u)^3.$$

From the small data global existence, we conclude that for the mass-critical Hartree equation (1), there exists a minimal mass m_0 such that solutions with mass strictly smaller than m_0 are global and scatter in time. It is conjectured that m_0 should be $+\infty$ in the defocusing case and be $M(Q)$ in the focusing case, where Q is the *ground state*, that is, the positive radial Schwartz solution Q to the elliptic equation

$$\Delta Q + (|x|^{-2} * |Q|^2)Q = Q. \quad (6)$$

In this paper we prove the conjecture for radial data. In particular, we have

Theorem 1.2. *In the defocusing case $\mu = +1$, all maximal-lifespan radial solutions to (1) are global and do not blow up either forward or backward in time. In the focusing case $\mu = -1$, all maximal-lifespan radial solutions to (1) with $M(u) < M(Q)$ are global and do not blow up either forward or backward in time.*

In fact, the result in Theorem 1.2 is sharp. $e^{it}Q$ is the solution to (1) that blows up at infinity. Moreover, this equation is invariant under the pseudo-conformal transformation

$$u(t, x) \mapsto (i(t - T))^{-\frac{d}{2}} e^{\frac{i|x|^2}{4(t-T)}} \bar{u}\left(\frac{1}{t-T}, \frac{x}{t-T}\right).$$

So $(i(t - T))^{-\frac{d}{2}} e^{\frac{i|x|^2}{4(t-T)}} e^{-\frac{i}{t-T}} Q\left(\frac{x}{t-T}\right)$ is the solution that blows up at finite time $t = T$ for fixed T .

In the proof of the above theorem, we adapt the ideas and techniques in [11] and [12], which represent the state of the art in nonlinear dispersive equations. In [11], R. Killip, T. Tao and M. Visan established the global well-posedness and scattering for radial solutions to mass-critical nonlinear Schrödinger equations in dimension $d = 2$. R. Killip, M. Visan and X. Zhang extended this result to higher dimensions in [12]. In addition, C. E. Kenig, F. Merle dealt with the focusing energy-critical nonlinear Schrödinger equation with radial data in [8]. For other related works, see S. Keraani [10], T. Tao, M. Visan and X. Zhang [22], [23]. Before we state our argument, we need some definitions.

Definition 1.3 (Symmetry group). *For any phase $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, position $x_0 \in \mathbb{R}^d$, frequency $\xi_0 \in \mathbb{R}^d$ and scaling parameter $\lambda > 0$, we define the unitary transformation $g_{\theta, \xi_0, x_0, \lambda} : L_x^2(\mathbb{R}^d) \rightarrow L_x^2(\mathbb{R}^d)$ by the formula*

$$[g_{\theta, \xi_0, x_0, \lambda} f](x) := \frac{1}{\lambda^{d/2}} e^{i\theta} e^{ix \cdot \xi_0} f\left(\frac{x - x_0}{\lambda}\right).$$

We let G be the collection of such transformations. We also let $G_{rad} \subset G$ denote the collection of transformations in G which preserve spherical symmetry, or more explicitly,

$$G_{rad} := \{g_{\theta, 0, 0, \lambda} : \theta \in \mathbb{R}/2\pi\mathbb{Z}; \lambda > 0\}.$$

Definition 1.4 (Almost periodicity modulo symmetries). *A solution u with lifespan I is said to be almost periodic modulo G if there exist (possibly discontinuous) functions $N : I \rightarrow \mathbb{R}^+$, $\xi : I \rightarrow \mathbb{R}^d$, $x : I \rightarrow \mathbb{R}^d$ and a function $C : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that*

$$\int_{|x-x(t)| \geq C(\eta)/N(t)} |u(t, x)|^2 dx \leq \eta$$

and

$$\int_{|\xi-\xi(t)| \geq C(\eta)N(t)} |\hat{u}(t, \xi)|^2 d\xi \leq \eta$$

for all $t \in I$ and $\eta > 0$. We refer to the function N as the frequency scale function for the solution u , ξ as the frequency center function, and C as the compactness modulus function. Furthermore, if we can select $x(t) = \xi(t) = 0$ for all $t \in I$, then we say that u is almost periodic modulo G_{rad} .

Remark 1.1. *By Ascoli-Arzelà theorem, the above definition is equivalent to either of the following two statements:*

- I. *The quotient orbit $\{Gu(t) : t \in I\}$ is a precompact set of $G \backslash L^2$, where $G \backslash L^2$ is the moduli space of G -orbits $Gf := \{gf : g \in G\}$ of $L^2(\mathbb{R}^d)$.*
- II. *There exists a compact subset K of L^2 such that $u(t) \in GK$ for all $t \in I$; equivalently there exists a group function $g : I \rightarrow G$ and a compact subset K such that $g^{-1}(t)u(t) \in K$ for any $t \in I$.*

Suppose for contradiction that Theorem 1.2 is not true, then we can find an almost periodic solution. The solution must be one of the following three forms:

Theorem 1.3 (Three special scenarios for blowup). *Suppose Theorem 1.2 failed for spherically symmetric solutions, then there exists a maximal-lifespan solution u which may be chosen to be spherically symmetric and almost periodic modulo G_{rad} . Moreover, it blows up both forward and backward in time, and in the focusing case also obeys $M(u) < M(Q)$.*

With spherical symmetry, we can also ensure that the lifespan I and the frequency scale function $N : I \rightarrow \mathbb{R}^+$ match one of the following three scenarios:

- I. (Soliton-like solution) *We have $I = \mathbb{R}$ and*

$$N(t) = 1$$

for all $t \in \mathbb{R}$.

- II. (Double high-to-low frequency cascade) *We have $I = \mathbb{R}$,*

$$\liminf_{t \rightarrow -\infty} N(t) = \liminf_{t \rightarrow +\infty} N(t) = 0,$$

and

$$\sup_{t \in \mathbb{R}} N(t) < \infty.$$

III. (Self-similar solution) We have $I = (0, +\infty)$ and

$$N(t) = t^{-1/2}$$

for all $t \in I$.

This is a wonderful classification theorem first given in [11] although some other authors have mentioned some of them, see [3], [6], [8], [20], [23], etc. In view of this theorem, our goal is to preclude the possibilities of all the scenarios.

Note that the minimal mass blow-up solution has very good properties because it is localized in both physical and frequency space. In fact, it admits higher regularity.

Theorem 1.4 (Regularity in the self-similar case). *Let u be a spherically symmetric solution to (1) that is almost periodic modulo G_{rad} and self-similar in the sense of Theorem 1.3. Then $u(t) \in H^s(\mathbb{R}^d)$ for all $t \in (0, \infty)$ and all $s \geq 0$.*

Theorem 1.5 (Regularity in the global case). *Let u be a global spherically symmetric solution to (1) that is almost periodic modulo G_{rad} . Suppose also that $N(t) \lesssim 1$ for all $t \in \mathbb{R}$, then $u \in L_t^\infty H^s(\mathbb{R} \times \mathbb{R}^d)$ for all $s \geq 0$.*

In the proofs of these two theorems for mass-critical Schrödinger equations in [11], the radial assumption is fully exploited based on a careful observation that there is a dichotomy between scattering solutions and almost periodic solutions. There are similar results for the mass-critical Hartree equation. More precisely, one has:

Proposition 1.1. *Let $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ be a maximal-lifespan solution which is almost periodic modulo G . Then $e^{-it\Delta}u(t)$ is weakly convergent to zero in $L_x^2(\mathbb{R}^d)$ as $t \rightarrow \sup(I)$ or $t \rightarrow \inf(I)$.*

As a corollary of Proposition 1.1, we have

Corollary 1.1 (A Duhamel formula). *Let u be a solution to (1) which is almost periodic modulo G . Its maximal-lifespan is I . Then for all $t \in I$,*

$$u(t) = \lim_{T \nearrow \sup I} i \int_t^T e^{i(t-t')\Delta} F(u(t')) dt' \quad (7)$$

$$= - \lim_{T \searrow \inf I} i \int_t^T e^{i(t-t')\Delta} F(u(t')) dt' \quad (8)$$

as weak limit in L_x^2 .

There are some new difficulties in dealing with the mass-critical Hartree equation. One of them comes from the asymptotic orthogonality. In the study of the mass-critical Hartree equation, we have to use the non-symmetric spacetime norm because the symmetric spacetime norm will lead to the restriction on dimension. However, the orthogonality can be destroyed by the non-symmetric spacetime norm.

We illustrate this by considering the simple example: Let φ_1 and φ_2 are two bump functions in $\mathbb{R} \times \mathbb{R}^2$. Let $x_n^1, x_n^2 \in \mathbb{R}^2$ be such that $|x_n^1 - x_n^2| \rightarrow \infty$ as $n \rightarrow \infty$. Then we have

$$\|\varphi_1(t, x + x_n^1) + \varphi_2(t, x + x_n^2)\|_{L_{t,x}^4}^4 \rightarrow \|\varphi_1\|_{L_{t,x}^4}^4 + \|\varphi_2\|_{L_{t,x}^4}^4, \quad (9)$$

as $n \rightarrow \infty$. However, if we replace $L_{t,x}^4$ with $L_t^6 L_x^3$, then

$$\begin{aligned} & \|\varphi_1(t, x + x_n^1) + \varphi_2(t, x + x_n^2)\|_{L_t^6 L_x^3}^6 \\ &= \int \left(\int |\varphi_1(t, x + x_n^1) + \varphi_2(t, x + x_n^2)|^3 dx \right)^2 dt \\ &= \int \left(\int (|\varphi_1(t, x + x_n^1)|^3 + |\varphi_2(t, x + x_n^2)|^3 \right. \\ & \quad \left. + 3|\varphi_1(t, x + x_n^1)|^2 |\varphi_2(t, x + x_n^2)| + 3|\varphi_1(t, x + x_n^1)| |\varphi_2(t, x + x_n^2)|^2) dx \right)^2 dt \\ &\rightarrow \int \left(\int (|\varphi_1(t, x + x_n^1)|^3 + |\varphi_2(t, x + x_n^2)|^3) dx \right)^2 dt \\ &= \int \left(\int |\varphi_1(t, x)|^3 dx \right)^2 dt + \int \left(\int |\varphi_2(t, x)|^3 dx \right)^2 dt \\ & \quad + 2 \int \left(\int |\varphi_1(t, x)|^3 dx \right) \left(\int |\varphi_2(t, x)|^3 dx \right) dt \\ &\rightarrow \|\varphi_1\|_{L_t^6 L_x^3}^6 + \|\varphi_2\|_{L_t^6 L_x^3}^6, \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Fortunately, for radial solution of the free Schrödinger equation, there are only two kinds of orthogonality - time translation and scaling (**NO** spatial translation!), both of which are possessed by time variable. So the orthogonality can be exploited and get the desired orthogonal relation similar to (9) (see Section 3). So the radial assumption is necessary to prove Theorem 1.3, which is in contrast to [11], where the similar theorem was established without the radial assumption. Such assumption is also used in precluding the three enemies in the sense of Theorem 1.3.

Some other difficulties coming from the convolution in the nonlinearity lie on the fact that it's non-local in physical space and singular in frequency space. For example, in precluding the self-similar solution, we need to deal with terms such as $(V * |u_{lo}^2|)u_{hi}$, where $\hat{u}_{lo}$ is supported in $\{\xi : |\xi| \leq M\}$ and $\hat{u}_{hi}$ is supported in $\{\xi : |\xi| \geq N\}$. In [11], the corresponding term $|u_{hi}|^2 u_{lo}$ can be estimated by means of bilinear estimate (Lemma 2.6). However, the convolution prevents the direct interaction between u_{lo} and u_{hi} in Hartree equation, so the bilinear estimate cannot be applied. In fact, to overcome the difficulty we exploit Shao's estimate (Lemma 2.5) and its dual estimate in full strength to replace the bilinear estimates (see Section 5). Meanwhile, we adapt weighted Strichartz estimate (16) in obtaining the additional regularity for the double high-to-low frequency cascade and soliton-like solutions, where the non-locality of the nonlinearity forces us to apply such estimates in different regions (see Section 6).

The rest of the paper is organized as follows: In Section 2, we record some known results such as basic facts in harmonic analysis, various versions of Strichartz estimates

and in/out decomposition. In Section 3, we give the stability theory and the concentration compactness result. In Section 4, we show that any failure of Theorem 1.2 must be “caused” by almost periodic solutions. In Section 5, we preclude the self-similar solution by proving that it possesses additional regularity. In Section 6, we prove the additional regularity in the other two cases. In Section 7 and Section 8, we preclude the double high-to-low frequency and soliton-like solutions.

2 Preliminaries

2.1 Some Notations

We use $X \lesssim Y$ or $Y \gtrsim X$ whenever $X \leq CY$ for some constant $C > 0$. We use $O(Y)$ to denote any quantity X such that $|X| \lesssim Y$. We use the notation $X \sim Y$ whenever $X \lesssim Y \lesssim X$. If C depends upon some additional parameters, we will indicate this with subscripts; for example, $X \lesssim_u Y$ denotes the assertion that $X \leq C_u Y$ for some C_u depending on u . We use the ‘Japanese bracket’ convention $\langle x \rangle = (1 + |x|^2)^{1/2}$. We write $L_t^q L_x^r$ to denote the Banach space with norm

$$\|u\|_{L_t^q L_x^r(\mathbb{R} \times \mathbb{R}^3)} := \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}^3} |u(t, x)|^r dx \right)^{q/r} dt \right)^{1/q},$$

with the usual modifications when q or r are equal to infinity, or when the domain $\mathbb{R} \times \mathbb{R}^d$ is replaced by spacetime slab such as $I \times \mathbb{R}^d$. When $q = r$ we abbreviate $L_t^q L_x^q$ as $L_{t,x}^q$.

2.2 Basic harmonic analysis

We recall some basic facts in Littlewood-Paley theory. Let $\varphi(\xi)$ be a radial bump function supported in the ball $\{\xi \in \mathbb{R}^d : |\xi| \leq \frac{11}{10}\}$ and equal to 1 on the ball $\{\xi \in \mathbb{R}^d : |\xi| \leq 1\}$. For each number $N > 0$, we define the Fourier multipliers

$$\begin{aligned} \widehat{P_{<N}} f(\xi) &:= \varphi(\xi/N) \hat{f}(\xi), \\ \widehat{P_{\geq N}} f(\xi) &:= (1 - \varphi(\xi/N)) \hat{f}(\xi), \\ \widehat{P_N} f(\xi) &:= (\varphi(\xi/N) - \varphi(2\xi/N)) \hat{f}(\xi) \end{aligned}$$

and similarly $P_{<N}$ and $P_{\geq N}$. We also define

$$P_{M < \cdot \leq N} := P_{\leq N} - P_{\leq M} = \sum_{M < N' \leq N} P_{N'}$$

whenever $M < N$. We will usually use these multipliers when M and N are dyadic numbers; in particular, all summations over N or M are understood to be over dyadic numbers. Nevertheless, it will occasionally be convenient to allow M and N to not

be a power of 2. Note that P_N is not truly a projection; to get around this, we will occasionally need to use fattened Littlewood-Paley operators:

$$\tilde{P}_N := P_{N/2} + P_N + P_{2N}. \quad (10)$$

They obey $P_N \tilde{P}_N = \tilde{P}_N P_N = P_N$.

As all Fourier multipliers, the Littlewood-Paley operators commute with the propagator $e^{it\Delta}$, as well as with differential operators such as $i\partial_t + \Delta$. We will use basic properties of these operators many times, including

Lemma 2.1 (Bernstein estimates). *For $1 \leq p \leq q \leq \infty$,*

$$\begin{aligned} \|\nabla^{\pm s} P_N f\|_{L_x^p(\mathbb{R}^d)} &\sim N^{\pm s} \|P_N f\|_{L_x^p(\mathbb{R}^d)}, \\ \|P_{\leq N} f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{p} - \frac{d}{q}} \|P_{\leq N} f\|_{L_x^p(\mathbb{R}^d)}, \\ \|P_N f\|_{L_x^q(\mathbb{R}^d)} &\lesssim N^{\frac{d}{p} - \frac{d}{q}} \|P_N f\|_{L_x^p(\mathbb{R}^d)}. \end{aligned}$$

2.3 Strichartz estimates

Naturally, everything that we do for Hartree equation builds on basic properties of the linear propagator $e^{it\Delta}$.

From the explicit formula

$$e^{it\Delta} f(x) = \frac{1}{(4\pi it)^{d/2}} \int_{\mathbb{R}^d} e^{i|x-y|^2/4t} f(y) dy,$$

we deduce the standard dispersive inequality

$$\|e^{it\Delta} f\|_{L^p(\mathbb{R}^d)} \lesssim \frac{1}{|t|^{d(1/2-1/p)}} \|f\|_{L^{p'}(\mathbb{R}^d)} \quad (11)$$

for all $t \neq 0$ and $2 \leq p \leq \infty$.

Lemma 2.2 (Kernel estimates, [11]). *For any $m \geq 0$, the kernel of the linear propagator obeys the following estimates:*

$$|(P_N e^{it\Delta})(x, y)| \lesssim_m \begin{cases} |t|^{-d/2} & : |x - y| \sim Nt \\ \frac{N^d}{|N^2 t|^m \langle N|x - y| \rangle^m} & : \text{otherwise} \end{cases} \quad (12)$$

for $|t| \geq N^{-2}$ and

$$|(P_N e^{it\Delta})(x, y)| \lesssim_m N^d \langle N|x - y| \rangle^{-m} \quad (13)$$

for $|t| \leq N^{-2}$.

Lemma 2.3 (Strichartz estimates, [21]). Fix $d \geq 1$ and call a pair (q, r) admissible if $2 \leq q, r \leq \infty$, $\frac{2}{q} + \frac{d}{r} = \frac{d}{2}$ and $(q, r, d) \neq (2, \infty, 2)$. Then for any admissible pair (q, r) and $(\tilde{q}, \tilde{r})$, let I be an interval, let $t_0 \in I$, and let $u_0 \in L_x^2(\mathbb{R}^d)$ and $f \in L_t^{\tilde{q}'} L_x^{\tilde{r}'}$. Then the function u defined by

$$u(t) := e^{i(t-t_0)\Delta} u_0 - i \int_{t_0}^t e^{i(t-t')\Delta} f(t') dt' \quad (14)$$

obeys the estimate

$$\|u\|_{L_t^q L_x^r} \lesssim \|u_0\|_{L_x^2} + \|f\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}} \quad (15)$$

where all spacetime norms are over $I \times \mathbb{R}^d$.

Lemma 2.4 (Weighted Strichartz, [12]). Let I be an interval, let $t_0 \in I$, $u_0 \in L_x^2(\mathbb{R}^d)$ and $f \in L_t^2 L_x^{\frac{2d}{d+2}}$ be spherically symmetric. Then the function u defined by (14) obeys the estimate

$$\left\| |x|^{\frac{2(d-1)}{q}} u \right\|_{L_t^q L_x^{\frac{2q}{q-4}}(I \times \mathbb{R}^d)} \lesssim \|u_0\|_{L_x^2(\mathbb{R}^d)} + \|f\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} \quad (16)$$

for all $4 \leq q \leq \infty$.

Lemma 2.5 (Shao's Strichartz estimate, [19]). Let $d \geq 2$, for $f \in L_{rad}^2(\mathbb{R}^d)$, we have

$$\|P_N e^{it\Delta} f\|_{L_{t,x}^q(\mathbb{R} \times \mathbb{R}^d)} \lesssim_q N^{\frac{d}{2} - \frac{d+2}{q}} \|f\|_{L_x^2(\mathbb{R}^d)}, \quad (17)$$

provided $q > \frac{4d+2}{2d-1}$.

Lemma 2.6 (Bilinear Strichartz, [4],[6]). For any spacetime slab $I \times \mathbb{R}^d$, any $t_0 \in I$ and any $M, N > 0$, we have

$$\begin{aligned} \|(P_{\geq N} u)(P_{\leq M} v)\|_{L_{t,x}^2(I \times \mathbb{R}^d)} &\lesssim_q \frac{M^{\frac{d-1}{2}}}{N^{-\frac{1}{2}}} \left(\|P_{\geq N} u(t_0)\|_{L^2(\mathbb{R}^d)} + \|(i\partial_t + \Delta) P_{\geq N} u\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} \right) \\ &\quad \times \left(\|P_{\leq M} v(t_0)\|_{L^2(\mathbb{R}^d)} + \|(i\partial_t + \Delta) P_{\leq M} v\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} \right) \end{aligned}$$

for all functions u, v on $I \times \mathbb{R}^d$.

2.4 An in/out decomposition

We define the projection onto outgoing spherical waves by

$$[P^+ f](r) = \frac{1}{2} \int_0^\infty r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(1)}(kr) \hat{f}(k) k^{\frac{d}{2}} dk$$

and the projection onto incoming spherical waves by

$$[P^- f](r) = \frac{1}{2} \int_0^\infty r^{\frac{2-d}{2}} H_{\frac{d-2}{2}}^{(2)}(kr) \hat{f}(k) k^{\frac{d}{2}} dk,$$

where $H_{\frac{d-2}{2}}^{(1)}$ denotes the Hankel function of the first kind with order $\frac{d-2}{2}$ and $H_{\frac{d-2}{2}}^{(2)}$ denotes the Hankel function of the second kind with order $\frac{d-2}{2}$. We will write $P_N^\pm$ for the product $P^\pm P_N$.

Lemma 2.7 (Kernel estimates, [12]). *For $|x| \gtrsim N^{-1}$ and $t \gtrsim N^{-2}$, the integral kernel obeys*

$$|[P_N^\pm e^{\mp it\Delta}](x, y)| \lesssim \begin{cases} (|x||y|)^{-\frac{d-1}{2}} |t|^{-\frac{1}{2}} & : |y| - |x| \sim Nt \\ \frac{N^d}{(N|x|)^{\frac{d-1}{2}} \langle N|y| \rangle^{\frac{d-1}{2}}} \langle N^2 t + N|x| - N|y| \rangle^{-m} & : \text{otherwise} \end{cases} \quad (18)$$

for any $m \geq 0$. For $|x| \gtrsim N^{-1}$ and $|t| \lesssim N^{-2}$, the integral kernel obeys

$$|[P_N^\pm e^{\mp it\Delta}](x, y)| \lesssim \frac{N^d}{(N|x|)^{\frac{d-1}{2}} \langle N|y| \rangle^{\frac{d-1}{2}}} \langle N|x| - N|y| \rangle^{-m}$$

for any $m \geq 0$.

Lemma 2.8 (Properties of $P^\pm$, [12]). *We have*

(i) $P^+ + P^-$ acts as the identity on $L_{rad}^2(\mathbb{R}^d)$.

(ii) Fix $N > 0$. For any spherically symmetric function $f \in L_x^2(\mathbb{R}^d)$,

$$\|P^\pm P_{\geq N} f\|_{L_x^2(|x| \geq \frac{1}{100} N^{-1})} \lesssim \|f\|_{L_x^2(\mathbb{R}^d)}$$

with an N -independent constant.

3 Stability and concentration compactness

Lemma 3.1 (Stability). *For every $A > 0$ and $\varepsilon > 0$, there exists $\delta > 0$ with the following property: if $u : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ approximately solves (1) in the sense that*

$$iu_t + \Delta u - F(u) = e,$$

with

$$\|e\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I \times \mathbb{R}^d)} \leq \delta$$

and obeys

$$\|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(I \times \mathbb{R}^d)} \leq A,$$

and $t_0 \in I$ and $v_0 \in L_x^2(\mathbb{R}^d)$ are such that

$$\|u(t_0) - v_0\|_{L_x^2} \leq \delta,$$

then there exists a solution $v : I \times \mathbb{R}^d \rightarrow \mathbb{C}$ to (1) with $v(t_0) = v_0$ such that

$$\|u - v\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(I \times \mathbb{R}^d)} \leq \varepsilon.$$

In particular, by the Strichartz inequality,

$$\|u - v\|_{L_t^\infty L_x^2(I \times \mathbb{R}^d)} \lesssim \delta + \varepsilon A^2.$$

Proof. We first establish this claim when A is sufficiently small depending on d . Let $v : I' \times \mathbb{R}^d \rightarrow \mathbb{C}$ be the maximal-lifespan solution with initial data $v(t_0)$. Writing $v = u + w$ on the interval $I'' := I \cap I'$, then w satisfies

$$iw_t + \Delta w = F(u + w) - F(u) - e$$

with

$$\|e^{i(t-t_0)\Delta} w(t_0)\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(I'' \times \mathbb{R}^d)} \leq C'_d \delta.$$

Let $X := \|w\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(I'' \times \mathbb{R}^d)}$, then by Lemma 2.3, we have

$$\begin{aligned} X &\leq C'_d \delta + C''_d \left(\|F(u + w) - F(u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(I'' \times \mathbb{R}^d)} + \delta \right) \\ &\leq \tilde{C}_d (A^2 X + A X^2 + X^3 + \delta) \end{aligned}$$

where $\tilde{C}_d$ depends only on d . If A is sufficiently small depending on d and δ is sufficiently small depending on ε and d , then the standard continuity argument gives $X \leq \varepsilon$. If A is large, we can iterate the case when A is small (shrinking δ , ε repeatedly) after a subdivision of the time interval. $\square$

We now need a key concentration-compactness result. The concentration compactness principle was first introduced by F. Merle, L. Vega [17] and Bahouri, P. Gerard [2] to study nonlinear Schrödinger equations. The idea was further developed by S. Keraani [10]. The results of [17] and [10] were extended to higher dimensions by P. Begout and A. Vargas [1]. Because the solution of the free Schrödinger equation is still a solution under the action of linear propagator $e^{it_0\Delta}$, we will need to enlarge the group G to contain this linear propagator.

Definition 3.1 (Enlarged group). *For any phase $\theta \in \mathbb{R}/2\pi\mathbb{Z}$, position $x_0 \in \mathbb{R}^d$, frequency $\xi_0 \in \mathbb{R}^d$, scaling parameter $\lambda > 0$, and time t_0 , we define the unitary transformation $g_{\theta, x_0, \xi_0, \lambda, t_0} : L_x^2(\mathbb{R}^d) \rightarrow L_x^2(\mathbb{R}^d)$ by the formula*

$$g_{\theta, x_0, \xi_0, \lambda, t_0} = g_{\theta, x_0, \xi_0, \lambda} e^{it_0\Delta}.$$

Let G' be the collection of such transformations. In particular, we denote by G'_{rad} the collection of all the transformation with $x_0 = \xi_0 = 0$. We also let G' act on global spacetime functions $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{C}$ by defining

$$T_{g_{\theta, \xi_0, x_0, \lambda, t_0}} u(t, x) := \frac{1}{\lambda^{\frac{d}{2}}} e^{i\theta} e^{ix \cdot \xi_0} e^{-it|\xi_0|^2} (e^{it_0\Delta} u) \left(\frac{t}{\lambda^2}, \frac{x - x_0 - 2\xi_0 t}{\lambda} \right).$$

Definition 3.2. For any two sequences g_n, g'_n in G' , we say that g_n and g'_n are asymptotically orthogonal if $(g_n)^{-1}g'_n$ diverges to infinity in G' . More explicitly, if $g_n = g_{\theta_n, \xi_n, x_n, \lambda_n, t_n}$ and $g'_n = g_{\theta'_n, \xi'_n, x'_n, \lambda'_n, t'_n}$, then this asymptotic orthogonality is equivalent to

$$\lim_{n \rightarrow \infty} \left(\frac{\lambda_n}{\lambda'_n} + \frac{\lambda'_n}{\lambda_n} + |t_n \lambda_n^2 - t'_n (\lambda'_n)^2| + |\xi_n - \xi'_n| + |x_n - x'_n| \right) = +\infty.$$

Careful computation shows that if g_n and g'_n are asymptotically orthogonal, then

$$\lim_{n \rightarrow \infty} \langle g_n f, g'_n f' \rangle_{L_x^2(\mathbb{R}^d)} = 0 \quad \text{for all } f, f' \in L_x^2(\mathbb{R}^d).$$

Theorem 3.1 (Linear profiles, [1]). *Fix d . Let $u_n, n = 1, 2, \dots$ be a bounded sequence in $L_{rad}^2(\mathbb{R}^d)$. Then (after passing to a subsequence if necessary) there exists a family $\phi^{(j)}$, $j = 1, 2, \dots$ of functions in $L_{rad}^2(\mathbb{R}^d)$ and group elements $g_n^{(j)} \in G'_{rad}$ for $j, n = 1, 2, \dots$ such that we have the decomposition*

$$u_n = \sum_{j=1}^l g_n^{(j)} \phi^{(j)} + w_n^{(l)} \quad (19)$$

for all $l = 1, 2, \dots$; here $w_n^l \in L_{rad}^2(\mathbb{R}^d)$ is such that its linear evolution has asymptotically vanishing scattering size:

$$\lim_{l \rightarrow \infty} \limsup_{n \rightarrow \infty} \|e^{it\Delta} w_n^l\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} = 0. \quad (20)$$

Moreover, $g_n^{(j)}$ and $g_n^{(j')} \in G'_{rad}$ are asymptotically orthogonal for any $j \neq j'$, and for any $l \geq 1$ we have the mass decoupling property

$$\lim_{n \rightarrow \infty} \left[M(u_n) - \sum_{j=1}^l M(\phi^{(j)}) - M(w_n^l) \right] = 0. \quad (21)$$

For later use, we prove the following lemma:

Lemma 3.2. *Let $g_n^{(j)}, g_n^{(j')} \in G'_{rad}$ be asymptotically orthogonal for any $j \neq j'$, then we have*

$$\left\| \sum_{j=1}^l g_n^{(j)} \phi^{(j)} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^6 \leq \sum_{j=1}^l \left\| g_n^{(j)} \phi^{(j)} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^6 + o(1), \quad \text{as } n \rightarrow \infty. \quad (22)$$

Proof. Since $d \geq 3$, $\frac{4}{3d-2} < 1$. So we have

$$\begin{aligned}
\left\| \sum_{j=1}^l g_n^{(j)} \phi^{(j)} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^6 &= \int \left(\int \left| \sum_{j=1}^l g_n^{(j)} \phi^{(j)} \right|^2 \left| \sum_{p=1}^l g_n^{(p)} \phi^{(p)} \right|^{\frac{4}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \\
&= \int \left(\sum_{j=1}^l \int \left| g_n^{(j)} \phi^{(j)} \right|^2 \left| \sum_{p=1}^l g_n^{(p)} \phi^{(p)} \right|^{\frac{4}{3d-2}} dx \right. \\
&\quad \left. + \sum_{j=1}^l \sum_{\substack{k=1 \\ k \neq j}}^l \int \left| g_n^{(j)} \phi^{(j)} g_n^{(k)} \phi^{(k)} \right| \left| \sum_{p=1}^l g_n^{(p)} \phi^{(p)} \right|^{\frac{4}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \\
&\leq \int \left(\sum_{j=1}^l \int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx + \sum_{j=1}^l \sum_{\substack{p=1 \\ p \neq j}}^l \int \left| g_n^{(j)} \phi^{(j)} \right|^2 \left| g_n^{(p)} \phi^{(p)} \right|^{\frac{4}{3d-2}} dx \right. \\
&\quad \left. + \sum_{j=1}^l \sum_{\substack{k=1 \\ k \neq j}}^l \int \left| g_n^{(j)} \phi^{(j)} g_n^{(k)} \phi^{(k)} \right| \left| \sum_{p=1}^l g_n^{(p)} \phi^{(p)} \right|^{\frac{4}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \\
&:= \int (A + B + C)^{\frac{3d-2}{d}} dt.
\end{aligned}$$

Without loss of generality, we can assume that all $\phi^{(j)}$ are compactly supported in both t and x . By orthogonality, B and C vanish as $n \rightarrow \infty$. Thus

$$\left\| \sum_{j=1}^l g_n^{(j)} \phi^{(j)} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^6 \leq \int \left(\sum_{j=1}^l \int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt + o(1).$$

Now we consider

$$\begin{aligned}
&\int \left(\sum_{j=1}^l \int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \\
&= \int \sum_{j=1}^l \left(\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^2 \left(\sum_{p=1}^l \int \left| g_n^{(p)} \phi^{(p)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{d-2}{d}} dt \\
&\quad + \sum_{j=1}^l \sum_{\substack{k=1 \\ k \neq j}}^l \int \left(\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right) \left(\int \left| g_n^{(k)} \phi^{(k)} \right|^{\frac{6d}{3d-2}} dx \right) \left(\sum_{p=1}^l \int \left| g_n^{(p)} \phi^{(p)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{d-2}{d}} dt \\
&:= I + II.
\end{aligned}$$

We estimate I first.

$$\begin{aligned}
I &\leq \sum_{j=1}^l \int \left(\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \\
&\quad + \sum_{j=1}^l \sum_{\substack{p=1 \\ p \neq j}}^l \int \left(\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^2 \left(\int \left| g_n^{(p)} \phi^{(p)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{d-2}{d}} dt. \tag{23}
\end{aligned}$$

Note that it does not change the compact support of time to take space norm, and

$$\begin{aligned}
\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx &= \frac{1}{(\rho_n^j)^{\frac{2d}{3d-2}}} \int \left| \phi^{(j)} \left(\frac{t}{(\rho_n^j)^2} - t_n^j, x \right) \right|^{\frac{6d}{3d-2}} dx \\
&:= \frac{1}{(\rho_n^j)^{\frac{2d}{3d-2}}} L^j \left(\frac{t}{(\rho_n^j)^2} - t_n^j \right),
\end{aligned}$$

we have

$$\begin{aligned}
&\int \left(\int \left| g_n^{(j)} \phi^{(j)} \right|^{\frac{6d}{3d-2}} dx \right)^2 \left(\int \left| g_n^{(k)} \phi^{(k)} \right|^{\frac{6d}{3d-2}} dx \right)^{\frac{d-2}{d}} dt \\
&= \left(\frac{\rho_n^j}{\rho_n^k} \right)^{\frac{2d-4}{3d-2}} \int |L^j(\tilde{t})|^2 \left| L^k \left(\left(\frac{\rho_n^j}{\rho_n^k} \right)^2 \tilde{t} + \left(\frac{\rho_n^j}{\rho_n^k} \right)^2 t_n^j - t_n^k \right) \right|^{\frac{d-2}{d}} d\tilde{t} \\
&\rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

We can prove similarly that $II \rightarrow 0$ when n is sufficiently large. $\square$

4 Almost periodic solutions

For brevity, we write $S_I(u)$ to denote $\|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(I \times \mathbb{R}^d)}$ in this section. If $I = \mathbb{R}$, we write $S_{\mathbb{R}}(u) = S(u)$.

Proposition 4.1. *Fix μ and d , and suppose that m_0 is finite. Let $u_n : I_n \times \mathbb{R}^d \rightarrow \mathbb{C}$ for $n = 1, 2, \dots$ be a sequence of radial solutions and $t_n \in I_n$ a sequence of times such that*

$$\limsup_{n \rightarrow \infty} M(u_n) = m_0,$$

and

$$\lim_{n \rightarrow \infty} S_{\geq t_n}(u_n) = \lim_{n \rightarrow \infty} S_{\leq t_n}(u_n) = \infty.$$

Then the sequence $G_{\text{rad}} u_n(t_n)$ has a subsequence which converges in $G_{\text{rad}} \setminus L_x^2$.

Proof. By time translation invariance, we may take $t_n = 0$ for all n . Then we have

$$\lim_{n \rightarrow \infty} S_{\geq 0}(u_n) = \lim_{n \rightarrow \infty} S_{\leq 0}(u_n) = \infty.$$

We consider the sequence of $\{u_n(0)\}$. Since $\limsup_{n \rightarrow \infty} M(u_n(0)) = m_0$, we have by concentration compactness principle that

$$u_n(0) = \sum_{j=1}^l g_n^{(j)} \varphi^{(j)} + w_n^l; \quad \text{with } g_n^{(j)} = h_n^{(j)} e^{it_n^j \Delta} \text{ and } h_n^{(j)} \in G_{rad}. \quad (24)$$

Moreover, we have the asymptotic orthogonality:

$$\|u_n(0)\|_{L_2}^2 = \sum_{j=1}^l \|\varphi^{(j)}\|_{L_2}^2 + \|w_n^l\|_{L_2}^2 + o(1), \quad (25)$$

and

$$\limsup_{n \rightarrow \infty} S(e^{it\Delta} w_n^l) \rightarrow 0 \quad \text{as } l \rightarrow \infty. \quad (26)$$

Claim: For any $\varepsilon > 0$, $\sup_j M(\varphi^{(j)}) \leq m_0 - \varepsilon$ doesn't hold.

Otherwise, there exists $\varepsilon_0 > 0$ such that $M(\varphi^{(j)}) \leq m_0 - \varepsilon_0$ for any $j = 1, 2, \dots$. Suppose $v^{(j)}$ is the nonlinear profile associated to $\varphi^{(j)}$ and depending on the limiting value of $t_n^{(j)}$, namely,

- ♣ If t_n^j is identically zero, $v^{(j)}$ is the maximal-lifespan solution to (1) with initial data $v^{(j)}(0) = \phi^{(j)}$.
- ♣ If t_n^j converges to $+\infty$, $v^{(j)}$ is the maximal-lifespan solution to (1) which scatters forward in time to $e^{it\Delta} \phi^{(j)}$.
- ♣ If t_n^j converges to $-\infty$, $v^{(j)}$ is the maximal-lifespan solution to (1) which scatters backward in time to $e^{it\Delta} \phi^{(j)}$.

Let

$$u_n^{(l)}(t) = \sum_{j=1}^l T_{h_n^{(j)}}[v^{(j)}(\cdot + t_n^j)](t) + e^{it\Delta} w_n^l, \quad (27)$$

then we have

$$\lim_{l \rightarrow +\infty} \lim_{n \rightarrow \infty} S(u_n^{(l)}) < +\infty. \quad (28)$$

In fact, by Lemma 3.2,

$$\lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} [S(u_n^{(l)})]^6 \leq \lim_{l \rightarrow \infty} \sum_{j=1}^l [S(v^{(j)})]^6.$$

Meanwhile, by (21), for any $\epsilon > 0$ sufficiently small, there exists j_0 such that

$$\|\varphi^{(j)}\|_{L_2}^2 < \epsilon, \quad \text{for all } j > j_0.$$

Note that $T_{h_n^j}$ preserves $L_t^6 L_x^{\frac{6d}{3d-2}}$ norm, by the small data theory, we conclude that for any $j > j_0$, the maximal lifespan $I^{(j)} = \mathbb{R}$ and

$$[S(v^{(j)})]^6 \leq CM(\varphi^{(j)})^3.$$

It follows that

$$\sum_{j > j_0} [S(v^{(j)})]^6 \leq \sum_{j > j_0} M(\varphi^{(j)})^3 \leq C,$$

where we use the mass decoupling property. For $j \leq j_0$, by the definition of m_0 and the fact that $\|v^{(j)}(0)\|_{L^2} \leq m_0 - \varepsilon_0$, we conclude that the maximal lifespan $I^{(j)} = \mathbb{R}$ and $S(v^{(j)}) \leq C$. Therefore, we have

$$\lim_{l \rightarrow +\infty} \lim_{n \rightarrow \infty} S(u_n^{(l)}) < \infty.$$

Meanwhile, by mass decoupling and the fact that $h_n^{(j)}$ preserves mass, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} M(u_n^{(l)}(0) - u_n(0)) &= \lim_{n \rightarrow \infty} M\left(\sum_{j=1}^l (T_{h_n^{(j)}}[v^{(j)}(\cdot + t_n^j)](0) - g_n^{(j)} \varphi^{(j)})\right) \\ &\leq \lim_{n \rightarrow \infty} \sum_{j=1}^l M(T_{h_n^{(j)}}[v^{(j)}(\cdot + t_n^j)](0) - g_n^{(j)} \varphi^{(j)}) \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^l M(h_n^{(j)}[v^{(j)}(t_n^j)] - h_n^{(j)} e^{it_n^j \Delta} \varphi^{(j)}) \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^l M(v^{(j)}(t_n^j) - e^{it_n^j \Delta} \varphi^{(j)}) = 0, \end{aligned}$$

where the last inequality follows from the definition of nonlinear profile.

Finally, we claim that

$$\lim_{l \rightarrow +\infty} \limsup_{n \rightarrow \infty} \left\| (i\partial_t + \Delta)u_n^{(l)} - F(u_n^{(l)}) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} = 0. \quad (29)$$

In fact, write

$$v_n^{(j)} := T_{h_n^{(j)}}[v^{(j)}(\cdot + t_n^j)].$$

By the definition of $u_n^{(l)}$, we have

$$u_n^{(l)} = \sum_{j=1}^l v_n^{(j)} + e^{it\Delta} w_n^l$$

and

$$(i\partial_t + \Delta)u_n^{(l)} = \sum_{j=1}^l F(v_n^{(j)}).$$

Moreover,

$$\begin{aligned} & \left\| (i\partial_t + \Delta)u_n^{(l)} - F(u_n^{(l)}) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \\ & \leq \left\| F(u_n^{(l)} - e^{it\Delta}w_n^{(l)}) - F(u_n^{(l)}) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} + \left\| \sum_{j=1}^l F(v_n^{(j)}) - F\left(\sum_{j=1}^l v_n^{(j)}\right) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}}. \end{aligned}$$

So it suffices to prove

$$\lim_{l \rightarrow +\infty} \limsup_{n \rightarrow \infty} \left\| F(u_n^{(l)} - e^{it\Delta}w_n^{(l)}) - F(u_n^{(l)}) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} = 0 \quad (30)$$

and

$$\lim_{n \rightarrow \infty} \left\| \sum_{j=1}^l F(v_n^{(j)}) - F\left(\sum_{j=1}^l v_n^{(j)}\right) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} = 0. \quad (31)$$

By Hardy-Littlewood-Sobolev inequality, we get

$$\begin{aligned} & \left\| F(u_n^{(l)} - e^{it\Delta}w_n^{(l)}) - F(u_n^{(l)}) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \\ & \lesssim \|e^{it\Delta}w_n^{(l)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} \|u_n^{(l)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^2 + \|e^{it\Delta}w_n^{(l)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^2 \|u_n^{(l)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} + \|e^{it\Delta}w_n^{(l)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^3. \end{aligned}$$

Thus (30) follows from (28) and (26). For (31), by Minkowski inequality, we have

$$\begin{aligned} & \left\| \sum_{j=1}^l F(v_n^{(j)}) - F\left(\sum_{j=1}^l v_n^{(j)}\right) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \\ & \lesssim \sum_{j_1 \neq j_2} \left\| (V * (v_n^{(j_1)} v_n^{(j_2)})) v_n^{(j_3)} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} + \sum_{j_1 \neq j_2} \left\| (V * |v_n^{(j_1)}|^2) v_n^{(j_2)} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}}. \end{aligned}$$

Since for any $j = 1, 2, \dots, l$, $v_n^{(j)}$ is the radial solution of (1) with data $\varphi^{(j)}$, $\|\varphi^{(j)}\|_{L^2} \leq m_0 - \varepsilon$, it follows from mass conservation and the definition of m_0 that

$$\|v_n^{(j)}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} \leq C. \quad (32)$$

Therefore, this together with orthogonality yields that

$$\begin{aligned} \left\| (V * (v_n^{(j_1)} v_n^{(j_2)})) v_n^{(j_3)} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} & \lesssim \left\| v_n^{(j_1)} v_n^{(j_2)} \right\|_{L_t^3 L_x^{\frac{3d}{3d-2}}} \left\| v_n^{(j_3)} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} \\ & \lesssim \left\| v_n^{(j_1)} v_n^{(j_2)} \right\|_{L_t^3 L_x^{\frac{3d}{3d-2}}} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned}$$

On the other hand, note that $v^{(j)}$ is radial, $h_n^{(j)} \in G_{rad}$ and that the orthogonality must be possessed by time variable, we have

$$\left\| (V * |v_n^{(j_1)}|^2) v_n^{(j_2)} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Thus, (31) follows. At last, by stability, we conclude that $\lim_{n \rightarrow \infty} S(u_n) < \infty$. This contradicts the hypothesis.

From the above claim, it follows that $l = 1$. So $u_n(0) = h_n e^{it_n \Delta} \varphi + w_n$ with $M(\varphi) = m_0$. Thus $M(w_n) \rightarrow 0$, which implies that $S(e^{it \Delta} w_n) \rightarrow 0$ as $n \rightarrow \infty$. Without loss of generality, we may take h_n to be identity. If $t_n \rightarrow 0$, then $u_n(0) \rightarrow \varphi$. Thus $G_{rad} u_n(0) \rightarrow G_{rad} \varphi$ in $G \setminus L_x^2$. It suffices to consider the case of $t_n \rightarrow \pm \infty$. We only consider the case of $t_n \rightarrow +\infty$, the other case is similar. Then we have

$$\begin{aligned} \lim_{n \rightarrow \infty} S_{\geq 0}(e^{it \Delta} u_n(0)) &= \lim_{n \rightarrow \infty} S_{\geq 0}(e^{it \Delta} h_n e^{it_n \Delta} \varphi) = \lim_{n \rightarrow \infty} S_{\geq 0}(T_{h_n} e^{it \Delta} e^{it_n \Delta} \varphi) \\ &= \lim_{n \rightarrow \infty} S_{\geq 0}(e^{i(t+t_n) \Delta} \varphi) = \lim_{n \rightarrow \infty} S_{\geq t_n}(e^{it \Delta} \varphi). \end{aligned}$$

Since $S(e^{it \Delta} \varphi) < +\infty$, we have

$$\lim_{n \rightarrow \infty} S_{\geq t_n}(e^{it \Delta} \varphi) = 0.$$

By stability again, we have $\lim_{n \rightarrow \infty} S_{\geq 0}(u_n) = 0$ and we reach a contradiction. $\square$

Proof of Theorem 1.3. We will only prove the first half of the theorem because the proof of the second half is identical with that of [11], which relies only the structure of group G_{rad} , pseudo-conformal invariance of (1) and is combinatorial.

Suppose Theorem 1.2 failed, then there exists a sequence of radial solutions u_n of (1) with $M(u_n) \leq m_0$ and $\lim_{n \rightarrow \infty} S(u_n) = +\infty$. Suppose u_n is maximal lifespan solutions, then there exists $t_n \in I_n$ such that $\lim_{n \rightarrow \infty} S_{\geq t_n}(u_n) = \lim_{n \rightarrow \infty} S_{\leq t_n}(u_n) = \infty$. By translation invariance, we may take t_n to be zero. From Proposition 4.1, it follows that there exists $u_0 \in L^2$ such that $G_{rad} u_n(0) \rightarrow G_{rad} u_0$, namely, $g_n u_n(0) \rightarrow u_0$ for some $g_n \in G_{rad}$. Without loss of generality, we may assume that g_n is identity. Thus $u_n(0) \rightarrow u_0$. Moreover, $M(u_0) \leq m_0$.

Let u be the radial solution of (1) with initial data u_0 , then u blows up both forward and backward in time. In fact, if u doesn't blow up forward in time, we have $S_{\geq 0}(u) < +\infty$. By stability, we have $\limsup_{n \rightarrow +\infty} S_{\geq 0}(u_n) < +\infty$. This contradicts the asymptotically blow-up. Similarly, we can prove that u blows up backward in time. By the definition of m_0 , we have $m_0 \leq M(u_0)$. Thus we have $M(u_0) = m_0$.

Now we consider any sequence $G_{rad} u(t'_n)$ for $t'_n \in I_n$. Since u blows up forward and backward in time, we have $S_{\geq t'_n}(u) = S_{\leq t'_n}(u) = \infty$. Then by Proposition 4.1, we have $G_{rad} u(t'_n) \rightarrow G_{rad} u_0$ (up to subsequence). Therefore, $\{G_{rad} u(t), t \in I\}$ is precompact in $G_{rad} \setminus L_x^2$.

Proposition 4.2 (Spacetime bound). *Let u be a non-zero solution to (1) with lifespan I , which is almost periodic modulo G with frequency scale function $N : I \rightarrow \mathbb{R}^+$. If J is any subinterval of I , then*

$$\int_J N(t)^2 dt \lesssim_u \int_J \left(\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \lesssim_u 1 + \int_J N(t)^2 dt. \quad (33)$$

Proof. We first prove that

$$\int_J \left(\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \lesssim_u 1 + \int_J N(t)^2 dt. \quad (34)$$

Let $0 < \eta < 1$ to be chosen momentarily and partition J into subintervals I_j so that

$$\eta/2 \leq \int_{I_j} N(t)^2 dt \leq \eta, \quad (35)$$

this requires at most $\eta^{-1} \times \text{RHS}(33)$ intervals. For each j , we may choose $t_j \in I_j$ so that

$$N(t_j)^2 |I_j| \leq 2\eta. \quad (36)$$

By Strichartz estimates, we have the following estimates on the spacetime slab $I_j \times \mathbb{R}^d$

$$\begin{aligned} \|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} &\lesssim \left\| e^{i(t-t_j)\Delta} u(t_j) \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} + \|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^3 \\ &\lesssim \|u_{>N_0}(t_j)\|_{L_x^2} + \left\| e^{i(t-t_j)\Delta} u_{<N_0}(t_j) \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}} + \|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^3 \\ &\lesssim \|u_{>N_0}(t_j)\|_{L_x^2} + |I_j|^{\frac{1}{6}} N_0^{\frac{1}{3}} \|u(t_j)\|_{L_x^2} + \|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^3 \\ &\lesssim \|u_{>N_0}(t_j)\|_{L_x^2} + \eta^{\frac{1}{6}} \|u(t_j)\|_{L_x^2} + \|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}}^3. \end{aligned}$$

Choosing N_0 as a large multiple of $N(t_j)$ and using Definition 1.4, one can make the first term arbitrary small. Choosing η sufficiently small depending on $M(u)$, one may also render the second term arbitrarily small. Thus by the bootstrap argument we obtain

$$\int_{I_j} \left(\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \lesssim \eta.$$

So (34) follows if we use the bound on the number of intervals I_j .

Now we prove

$$\int_J \left(\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \right)^{\frac{3d-2}{d}} dt \gtrsim_u \int_J N(t)^2 dt. \quad (37)$$

Using Definition 1.4 and choosing η sufficiently small depending on $M(u)$, we can guarantee that

$$\int_{|x-x(t)| \leq C(\eta)N(t)^{-1}} |u(t, x)|^2 dx \gtrsim_u 1. \quad (38)$$

By Hölder inequality, we get

$$\int_{\mathbb{R}^d} |u(t, x)|^{\frac{6d}{3d-2}} dx \gtrsim \left(\int_{|x-x(t)| \leq C(\eta)N(t)^{-1}} |u(t, x)|^2 dx \right)^{\frac{3d}{3d-2}} N(t)^{\frac{2d}{3d-2}}.$$

Using (38) and integrating over J we derive (37). $\square$

5 The self-similar solutions

This section is devoted to proving Theorem 1.4. Let u be as in Theorem 1.4. For any $A > 0$, we define

$$\mathcal{M}(A) := \sup_{T>0} \|u_{>AT^{-1/2}}(T)\|_{L_x^2(\mathbb{R}^d)}, \quad (39)$$

$$\mathcal{S}(A) := \sup_{T>0} \|u_{>AT^{-1/2}}(T)\|_{L_t^6 L_x^{\frac{6d}{3d-2}}([T, 2T] \times \mathbb{R}^d)}, \quad (40)$$

$$\mathcal{N}(A) := \sup_{T>0} \|P_{>AT^{-1/2}}F(u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)}, \quad (41)$$

where $u_{>AT^{-1/2}}(T) = P_{>AT^{-1/2}}u(T)$. To prove Theorem 1.4, it suffices to show that for every $s > 0$,

$$\mathcal{M}(A) \lesssim_{s,u} A^{-s}$$

whenever A is sufficiently large depending on u and s . From mass conservation, Proposition 4.2, self-similarity and Hardy-Littlewood-Sobolev inequality, we have

$$\mathcal{M}(A) + \mathcal{S}(A) + \mathcal{N}(A) \lesssim_u 1 \quad (42)$$

for all $A > 0$. From Strichartz estimates, we also see that

$$\mathcal{S}(A) \lesssim \mathcal{M}(A) + \mathcal{N}(A) \quad (43)$$

for all $A > 0$. A similar application of Strichartz estimates shows that for any admissible pair (q, r) ,

$$\|u\|_{L_t^q L_x^r([T, 2T] \times \mathbb{R}^d)} \lesssim_u 1 \quad (44)$$

for all $T > 0$.

Lemma 5.1 (Nonlinear estimate). *For all $A > 100$, we have*

$$\mathcal{N}(A) \lesssim_u \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}) + A^{-\frac{1}{2d+6}} [\mathcal{M}\left(\frac{A}{8}\right) + \mathcal{N}\left(\frac{A}{8}\right)].$$

Proof. It suffices to prove that

$$\left\| P_{>AT^{-\frac{1}{2}}}(F(u)) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \lesssim_u \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}) + A^{-\frac{1}{2d+6}} [\mathcal{M}\left(\frac{A}{8}\right) + \mathcal{N}\left(\frac{A}{8}\right)] \quad (45)$$

for arbitrary $T > 0$. To do this, we decompose u as

$$u = u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} + u_{\frac{1}{8}AT^{-\frac{1}{2}} < \cdot \leq \sqrt{A}AT^{-\frac{1}{2}}} + u_{< \sqrt{A}AT^{-\frac{1}{2}}}.$$

Then any term in the resulting expansion of $P_{>AT^{-\frac{1}{2}}}(F(u))$ that does not contain the factor of $u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}}$ vanishes.

Consider the terms which contain at least one factor of $u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}}$. The term which contains three factors of $u_{>\frac{1}{8}AT^{-\frac{1}{2}}}$ or which contains two factors of $u_{>\frac{1}{8}AT^{-\frac{1}{2}}}$ and one factor of $u_{<\sqrt{AT}^{-\frac{1}{2}}}$ can be estimated similarly. By Hölder inequality, Hardy-Littlewood-Sobolev inequality, (39), (40) and (44), we have

$$\begin{aligned}
& \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}} \right) u \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}} \right) \right\|_{L_t^6 L_x^{\frac{3d}{5}}([T, 2T] \times \mathbb{R}^d)} \|u\|_{L_t^3 L_x^{\frac{6d}{3d-4}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}} \right\|_{L_t^6 L_x^{\frac{3d}{3d-1}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \|u_{>\frac{1}{8}AT^{-\frac{1}{2}}}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}([T, 2T] \times \mathbb{R}^d)} \|u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}}\|_{L_t^\infty L_x^2([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}),
\end{aligned}$$

and

$$\begin{aligned}
& \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u \right) u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u \right) \right\|_{L_t^2 L_x^d([T, 2T] \times \mathbb{R}^d)} \|u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}}\|_{L_t^\infty L_x^2([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u \right\|_{L_t^2 L_x^{\frac{d}{d-1}}([T, 2T] \times \mathbb{R}^d)} \mathcal{M}(\sqrt{A}) \\
& \lesssim \|u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}}\|_{L_t^6 L_x^{\frac{6d}{3d-2}}([T, 2T] \times \mathbb{R}^d)} \|u\|_{L_t^3 L_x^{\frac{6d}{3d-4}}([T, 2T] \times \mathbb{R}^d)} \mathcal{M}(\sqrt{A}) \\
& \lesssim \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}).
\end{aligned}$$

Similarly, we have

$$\left\| V * \left(u_{\frac{1}{8}AT^{-\frac{1}{2}} > \cdot \geq \sqrt{AT}^{-\frac{1}{2}}} u \right) u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \lesssim \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}).$$

Finally, we consider the terms with one factor of $u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}}$ and two factors of $u_{<\sqrt{AT}^{-\frac{1}{2}}}$. First, by Hölder inequality, Lemma 2.1, Lemma 2.6, (39), (41) and (44),

we have

$$\begin{aligned}
& \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u_{< \sqrt{AT}^{-\frac{1}{2}}} \right) u_{< \sqrt{AT}^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| V * \left(u_{\geq \frac{1}{8}AT^{-\frac{1}{2}}} u_{< \sqrt{AT}^{-\frac{1}{2}}} \right) \right\|_{L_t^2 L_x^d([T, 2T] \times \mathbb{R}^d)} \left\| u_{< \sqrt{AT}^{-\frac{1}{2}}} \right\|_{L_t^\infty L_x^2([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim u \frac{1}{\left(\frac{1}{8}AT^{-\frac{1}{2}}\right)^{\frac{d-2}{2}}} \left\| u_{> \frac{1}{8}AT^{-\frac{1}{2}}} u_{< \sqrt{AT}^{-\frac{1}{2}}} \right\|_{L_{t,x}^2} \\
& \lesssim u A^{-\frac{d-1}{4}} \left(\left\| P_{> \frac{1}{8}AT^{-\frac{1}{2}}} u(t_0) \right\|_{L_x^2} + \left\| (i\partial_t + \Delta) P_{> \frac{1}{8}AT^{-\frac{1}{2}}} u \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \right) \\
& \quad \times \left(\left\| P_{< \sqrt{AT}^{-\frac{1}{2}}} u(t_0) \right\|_{L_x^2} + \left\| (i\partial_t + \Delta) P_{< \sqrt{AT}^{-\frac{1}{2}}} u \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}} \right) \\
& \lesssim u A^{-\frac{d-1}{4}} \left(\mathcal{M}\left(\frac{A}{8}\right) + \mathcal{N}\left(\frac{A}{8}\right) \right).
\end{aligned}$$

Now it suffices to estimate

$$\left\| \left(V * |u_{< \sqrt{AT}^{-\frac{1}{2}}}|^2 \right) u_{> \frac{1}{8}AT^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)}.$$

We divide it into two terms.

$$\begin{aligned}
& \left\| \left(V * |u_{< \sqrt{AT}^{-\frac{1}{2}}}|^2 \right) u_{> \frac{1}{8}AT^{-\frac{1}{2}}} \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& \leq \left\| \left(V * |u_{< \sqrt{AT}^{-\frac{1}{2}}}|^2 \right) P_{> \frac{1}{8}AT^{-\frac{1}{2}}} e^{i(t-T)\Delta} u(T) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& \quad + \left\| \left(V * |u_{< \sqrt{AT}^{-\frac{1}{2}}}|^2 \right) \int_T^t P_{> \frac{1}{8}AT^{-\frac{1}{2}}} e^{i(t-t')\Delta} F(u(t')) dt' \right\|_{L_t^2 L_x^{\frac{2d}{d+2}}([T, 2T] \times \mathbb{R}^d)} \\
& := I + II.
\end{aligned}$$

Let

$$q = \frac{2d+6}{d+1}, \quad s = -\frac{d}{2} + \frac{d+2}{q} = \frac{1}{d+3} > 0$$

and q_1, r_1, r_2 satisfy

$$\frac{1}{q_1} + \frac{1}{q} = \frac{1}{2}, \quad \frac{1}{r_1} + \frac{1}{q} = \frac{d+2}{2d}, \quad \frac{1}{r_1} = \frac{1}{r_2} - \frac{d-2+s}{d},$$

which yields that

$$\frac{2}{2q_1} + \frac{d}{2r_2} = \frac{d}{2}.$$

Thus, By Hölder inequality, Sobolev imbedding, Lemma 2.1, Lemma 2.5 and (44), we

have

$$\begin{aligned}
I &\leq \|V * |u|_{<\sqrt{AT}^{-\frac{1}{2}}}^2\|_{L_t^{q_1} L_x^{r_1}([T, 2T] \times \mathbb{R}^d)} \|P_{>\frac{1}{8}AT^{-\frac{1}{2}}} e^{i(t-T)\Delta} u(T)\|_{L_{t,x}^q([T, 2T] \times \mathbb{R}^d)} \\
&\leq C \| |\nabla|^s (|u|_{<\sqrt{AT}^{-\frac{1}{2}}}^2) \|_{L_t^{q_1} L_x^{r_2}([T, 2T] \times \mathbb{R}^d)} \sum_{N > \frac{1}{8}AT^{-\frac{1}{2}}} \|P_N e^{i(t-T)\Delta} u_{>\frac{1}{8}AT^{-\frac{1}{2}}}(T)\|_{L_{t,x}^q([T, 2T] \times \mathbb{R}^d)} \\
&\lesssim (\sqrt{AT}^{-\frac{1}{2}})^s \|u\|_{L_t^{2q_1} L_x^{2r_2}([T, 2T] \times \mathbb{R}^d)}^2 \sum_{N > \frac{1}{8}AT^{-\frac{1}{2}}} N^{\frac{d}{2} - \frac{d+2}{q}} \|P_{>\frac{1}{8}AT^{-\frac{1}{2}}} u(T)\|_{L_x^2} \\
&\lesssim_u (\sqrt{AT}^{-\frac{1}{2}})^s (AT^{-\frac{1}{2}})^{\frac{d}{2} - \frac{d+2}{q}} \mathcal{M}\left(\frac{A}{8}\right) \lesssim_u A^{-\frac{1}{2d+6}} \mathcal{M}\left(\frac{A}{8}\right).
\end{aligned}$$

Now we estimate II . By duality and Lemma 2.5, we get that

$$\left\| \int_{\mathbb{R}} P_N e^{-it\Delta} f(t, x) dt \right\|_{L_x^2} \leq C N^{\frac{d}{2} - \frac{d+2}{q}} \|f\|_{L_{t,x}^{q'}}. \quad (46)$$

Therefore, this together with Lemma 2.3 yields that for $q > \frac{4d+2}{2d-1}$,

$$\begin{aligned}
&\left\| \int_{\mathbb{R}} P_N e^{i(t-t')\Delta} F(t') dt' \right\|_{L_{t,x}^q} \\
&= \sup_{\|g\|_{L_{t,x}^{q'}}=1} \int_{\mathbb{R}} \left\langle \int_{\mathbb{R}} P_N e^{i(t-t')\Delta} F(t') dt', g(t, x) \right\rangle dt \\
&\leq \sup_{\|g\|_{L_{t,x}^{q'}}=1} \left\| \int_{\mathbb{R}} e^{-it'\Delta} F(t', x) dt' \right\|_{L_x^2} \left\| \int_{\mathbb{R}} P_N e^{-it\Delta} g(t, x) dt \right\|_{L_x^2} \\
&\lesssim N^{\frac{d}{2} - \frac{d+2}{q}} \|F\|_{L_t^2 L_x^{\frac{2d}{d+2}}}.
\end{aligned}$$

From this and Christ-Kiselev lemma (see [21]), we get that

$$\left\| \int_{t_0}^t P_N e^{i(t-t')\Delta} P_{>\frac{1}{8}AT^{-\frac{1}{2}}} F(u(t')) dt' \right\|_{L_{t,x}^q} \leq C N^{\frac{d}{2} - \frac{2d}{d+2}} \|P_{>\frac{1}{8}AT^{-\frac{1}{2}}} F\|_{L_t^2 L_x^{\frac{2d}{d+2}}}.$$

Similar to the estimate of I , we get that

$$II \lesssim_u A^{-\frac{1}{2d+6}} \mathcal{N}\left(\frac{A}{8}\right).$$

□

Lemma 5.2 (Qualitative decay). *We have*

$$\lim_{A \rightarrow \infty} \mathcal{M}(A) = \lim_{A \rightarrow \infty} \mathcal{S}(A) = \lim_{A \rightarrow \infty} \mathcal{N}(A) = 0. \quad (47)$$

Proof. Since u is almost periodic modulo G_{rad} and self-similar, for any $\eta > 0$, there exists $C(\eta)$, such that

$$\int_{|\xi| > C(\eta)T^{-\frac{1}{2}}} |\hat{u}(T, \xi)|^2 d\xi < \eta.$$

In particular, when A is sufficiently large, we have

$$\int_{|\xi| > AT^{-\frac{1}{2}}} |\hat{u}(T, \xi)|^2 d\xi < \eta.$$

Therefore, we get

$$\lim_{A \rightarrow \infty} \mathcal{M}(A) = 0.$$

From Lemma 5.1 and (42), we have

$$\begin{aligned} \mathcal{N}(A) &\leq \mathcal{S}\left(\frac{A}{8}\right) \mathcal{M}(\sqrt{A}) + A^{-\frac{1}{2d+6}} \left(\mathcal{M}\left(\frac{A}{8}\right) + \mathcal{N}\left(\frac{A}{8}\right)\right) \\ &\lesssim_u \mathcal{M}(\sqrt{A}) + A^{-\frac{1}{2d+6}} \rightarrow 0 \quad \text{as } A \rightarrow \infty. \end{aligned}$$

Finally, (43) yields

$$\mathcal{S}(A) \lesssim \mathcal{M}(A) + \mathcal{N}(A) \rightarrow 0 \quad \text{as } A \rightarrow \infty.$$

□

Proposition 5.1. *Let $0 < \eta < 1$. Then if A is sufficiently large depending on u and η ,*

$$\mathcal{S}(A) \leq \eta \mathcal{S}\left(\frac{A}{16}\right) + A^{-\frac{1}{2d+6}}. \quad (48)$$

In particular, $\mathcal{S}(A) \lesssim_u A^{-\frac{1}{2d+6}}$ for all $A > 0$.

Proof. Fix $\eta \in (0, 1)$. It suffices to show that

$$\left\| u_{>AT^{-\frac{1}{2}}} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}} [T, 2T] \times \mathbb{R}^d} \leq \eta \mathcal{S}\left(\frac{A}{16}\right) + A^{-\frac{1}{2d+6}} \quad (49)$$

for all $T > 0$.

Fix $T > 0$. By the Duhamel formula and then using Lemma 2.3, we obtain

$$\begin{aligned} \left\| u_{>AT^{-\frac{1}{2}}} \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}} [T, 2T] \times \mathbb{R}^d} &\lesssim \left\| P_{>AT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}} [T, 2T] \times \mathbb{R}^d} \\ &\quad + \left\| P_{>AT^{-\frac{1}{2}}} F(u) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}} ([\frac{T}{2}, 2T] \times \mathbb{R}^d)}. \end{aligned}$$

First, we consider the second term. By definition, we have

$$\left\| P_{>AT^{-\frac{1}{2}}} F(u) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}} ([\frac{T}{2}, 2T] \times \mathbb{R}^d)} \lesssim \mathcal{N}(A/2).$$

Using Lemma 5.1 and Lemma 5.2, we derive that

$$\left\| P_{>AT^{-\frac{1}{2}}} F(u) \right\|_{L_t^2 L_x^{\frac{2d}{d+2}} ([\frac{T}{2}, 2T] \times \mathbb{R}^d)} \lesssim RHS(48).$$

Thus the second term is acceptable.




Figure 1. Interpolation game board.

Now we consider the first term (see Figure 1). In fact, we will show that

$$\left\| P_{>AT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}[T, 2T] \times \mathbb{R}^d} \lesssim_u A^{-\frac{2(d-1)(d-2)}{3(4d^2-5d-2)}}, \quad (50)$$

which is acceptable, since $\frac{2(d-1)(d-2)}{3(4d^2-5d-2)} > \frac{1}{2d+6}$ for all $d \geq 3$. From Shao's Strichartz estimate (17), we have

$$\left\| P_M e^{it\Delta} f \right\|_{L_{t,x}^{\frac{2d}{d-1}}} \leq CM^{-\frac{d-2}{2d}} \|f\|_{L_x^2}. \quad (51)$$

Meanwhile, the Strichartz estimate gives

$$\|e^{it\Delta} f\|_{L_t^\infty L_x^2} \leq C \|f\|_{L_x^2}. \quad (52)$$

Interpolating between (51) and (52) with $\theta = 2/3$, we have

$$\left\| P_M e^{it\Delta} f \right\|_{L_t^{\frac{3d}{d-1}} L_x^{\frac{6d}{3d-2}}} \leq CM^{-\frac{d-2}{3d}} \|f\|_{L_x^2}. \quad (53)$$

Thus, from the mass conservation, we have

$$\left\| P_{BT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) \right\|_{L_t^{\frac{3d}{d-1}} L_x^{\frac{6d}{3d-2}}} \lesssim_u (BT^{-\frac{1}{2}})^{-\frac{d-2}{3d}}. \quad (54)$$

Using the Duhamel formula, we write

$$P_{BT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) = P_{BT^{-\frac{1}{2}}} e^{i(t-\varepsilon)\Delta} u(\varepsilon) - i \int_{\varepsilon}^{\frac{T}{2}} P_{BT^{-\frac{1}{2}}} e^{i(t-t')\Delta} F(u(t')) dt'$$

for any $\varepsilon > 0$. By self-similarity, the first term of RHS converges strongly to zero in L_x^2 as $\varepsilon \rightarrow 0$. By Lemma 2.1, it also converges to zero in $L_x^{\frac{6d}{3d-2}}$. Thus using the dispersive

estimate (11), we obtain

$$\begin{aligned}
& \left\| P_{BT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) \right\|_{L_t^{4d+2} L_x^{\frac{6d}{3d-2}}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim \left\| \int_0^{\frac{T}{2}} \frac{1}{|t-t'|^{\frac{1}{3}}} \|F(u(t'))\|_{L_x^{\frac{6d}{3d+2}}} dt' \right\|_{L_t^{4d+2}([T, 2T] \times \mathbb{R}^d)} \\
& \lesssim T^{\frac{1}{4d+2}-\frac{1}{3}} \sum_{0 < \tau < \frac{T}{4}} \|F(u)\|_{L_t^1 L_x^{\frac{6d}{3d+2}}([\tau, 2\tau] \times \mathbb{R}^d)} \\
& \lesssim T^{\frac{1}{4d+2}-\frac{1}{3}} \sum_{0 < \tau < \frac{T}{4}} \tau^{\frac{1}{6}} \|u\|_{L_t^{\frac{18}{5}} L_x^{\frac{18d}{9d-10}}([\tau, 2\tau] \times \mathbb{R}^d)}^3 \\
& \lesssim_u T^{\frac{1}{4d+2}-\frac{1}{6}},
\end{aligned}$$

where the last inequality comes from (44). Interpolating between this estimate and (54) with $\theta = \frac{2d(d-1)}{4d^2-5d-2}$, we obtain that

$$\left\| P_{BT^{-\frac{1}{2}}} e^{i(t-\frac{T}{2})\Delta} u\left(\frac{T}{2}\right) \right\|_{L_t^6 L_x^{\frac{6d}{3d-2}}([T, 2T] \times \mathbb{R}^d)} \lesssim_u B^{-\frac{2(d-1)(d-2)}{3(4d^2-5d-2)}}.$$

Summing this over dyadic $B \geq A$ yields (50) and (48).

Finally, we explain why (48) implies $\mathcal{S}(A) \lesssim_u A^{-\frac{1}{2d+6}}$. Choosing $\eta = \frac{1}{2}$. Then there exists A_0 depending on u , so that (48) holds for all $A \geq A_0$. By (42), we need only bound $S(A)$ for $A \geq A_0$.

Choose $k \geq 1$ so that $2^{-4k}A \leq A_0 \leq 2^{-4(k-1)}A$. By iterating (48) k times and using (42),

$$\begin{aligned}
S(A) & \leq [1 + \eta 2^{\frac{4}{2d+6}} + \dots + (\eta 2^{\frac{4}{2d+6}})^k] A^{-\frac{1}{2d+6}} + \eta^k S(2^{-4k}A) \\
& \lesssim_u A^{-\frac{1}{2d+6}} + \eta^k \lesssim_u A^{-\frac{1}{2d+6}}.
\end{aligned}$$

Note that the last inequality uses the way we choose η and k . □

Corollary 5.1. *For any $A > 0$, we have*

$$\mathcal{M}(A) + \mathcal{N}(A) + \mathcal{S}(A) \lesssim_u A^{-\frac{1}{2d+6}}.$$

Proof. The bound on $\mathcal{S}(A)$ was derived in Proposition 5.1. This together with Lemma 5.1 and (42) yields the bound on $\mathcal{N}(A)$.

We now turn to the bound on $\mathcal{M}(A)$. By Corollary 1.1,

$$\|P_{>AT^{-1/2}} u(T)\|_{L_x^2} \lesssim \sum_{k=0}^{\infty} \left\| \int_{2^k T}^{2^{k+1}T} e^{i(T-t')\Delta} P_{>AT^{-1/2}} F(u(t')) dt' \right\|_{L_x^2}, \quad (55)$$

where weak convergence has become strong convergence because of the frequency projection and the fact that $N(t) = t^{-1/2} \rightarrow 0$ as $t \rightarrow \infty$. Combining (55) with Strichartz estimates, (39) and (41), we get

$$\mathcal{M}(A) = \sup_{T>0} \|P_{>AT^{-1/2}} u(T)\|_{L_x^2} \lesssim \sum_{k=0}^{\infty} \mathcal{N}(2^{k/2} A). \quad (56)$$

The desired bound on $\mathcal{M}(A)$ now follows from that on $\mathcal{N}(A)$. $\square$

Proof of Theorem 1.4. Combining Lemma 5.1 with Corollary 5.1, one has

$$\mathcal{N}(A) \lesssim_u A^{-\frac{1}{4d+12}} [\mathcal{S}(\frac{A}{8}) + \mathcal{M}(\frac{A}{8}) + \mathcal{N}(\frac{A}{8})].$$

Together with (55) and (56), this allows us to deduce

$$\mathcal{S}(A) + \mathcal{M}(A) + \mathcal{N}(A) \lesssim_u A^{-\sigma} \quad \Rightarrow \quad \mathcal{S}(A) + \mathcal{M}(A) + \mathcal{N}(A) \lesssim_u A^{-\sigma - \frac{1}{4d+12}}$$

for any $\sigma > 0$. Iterating this statement shows that $u(t) \in H_x^s(\mathbb{R}^d)$ for all $s > 0$.

Corollary 5.2 (Absence of self-similar solutions). *There are no non-zero spherically symmetric solutions to (1) that are self-similar in the sense of Theorem 1.3.*

Proof. By Theorem 1.4, any such solution would obey $u(t) \in H_x^1(\mathbb{R}^d)$ for all $t \in (0, \infty)$. Then there exists a global solution with initial data $u(t_0)$ at any time $t_0 \in (0, \infty)$; recall that we assume $M(u) < M(Q)$ in the focusing case (see [13], [16]). On the other hand, self-similar solutions blow up at time $t = 0$. These two facts yield a contradiction. $\square$

6 Additional regularities

This section is devoted to the proof of Theorem 1.5. Before giving the proof, we record some basic local estimates. From mass conservation we have

$$\|u\|_{L_t^\infty L_x^2} \lesssim_u 1, \quad (57)$$

while from Definition 1.4 and the fact that $N(t)$ is bounded, we have

$$\lim_{N \rightarrow \infty} \|u_{\geq N}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} = 0. \quad (58)$$

From Proposition 4.2 and $N(t) \lesssim 1$, we have

$$\|u\|_{L_t^6 L_x^{\frac{6d}{3d-2}}(J \times \mathbb{R}^d)} \lesssim_u \langle |J| \rangle^{\frac{1}{6}} \quad (59)$$

for all intervals $J \subset \mathbb{R}$. By Hölder's inequality and Hardy-Littlewood-Sobolev inequality, this implies

$$\|F(u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} \lesssim_u \langle |J| \rangle^{\frac{1}{2}} \quad (60)$$

and then by Strichartz estimates (Lemma 2.3),

$$\|u\|_{L_t^q L_x^r(J \times \mathbb{R}^d)} \lesssim \langle |J| \rangle^{\frac{1}{q}}, \quad (61)$$

for any admissible pair (q, r) . Similarly, the weighted Strichartz estimates imply that

$$\| |x|^{\frac{2(d-1)}{q}} u \|_{L_t^q L_x^{\frac{2q}{q-4}}(J \times \mathbb{R}^d)} \lesssim_u \langle |J| \rangle^{\frac{1}{q}}. \quad (62)$$

Now for any dyadic number N , define

$$\mathcal{M}(N) := \|u_{\geq N}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)}, \quad (63)$$

then we see that $\mathcal{M}(N) \lesssim_u 1$ and

$$\lim_{N \rightarrow \infty} \mathcal{M}(N) = 0. \quad (64)$$

To prove Theorem 1.5, it suffices to show that $\mathcal{M}(N) \lesssim_{u,s} N^{-s}$ for any $s > 0$ and all N sufficiently large depending on u and s . This will immediately follow from iterating the following proposition with a suitably choice of small η (depending on u and s):

Proposition 6.1 (Regularity). *Let u be as in Theorem 1.5 and let $\eta > 0$ be a small number. Then*

$$\mathcal{M}(N) \lesssim_u \eta \mathcal{M}\left(\frac{N}{64}\right) \quad (65)$$

whenever N is sufficiently large depending on u and η .

The rest of this section is devoted to proving Proposition 6.1. Our task is to show that

$$\|u_{\geq N}(t_0)\|_{L_x^2(\mathbb{R}^d)} \lesssim_u \eta \mathcal{M}\left(\frac{N}{64}\right)$$

for all times t_0 and all N sufficiently large (depending on u and η). By time translation symmetry, we may assume $t_0 = 0$. By Corollary 1.1, we have

$$\begin{aligned} u_{\geq N}(0) &= (P^+ + P^-)u_{\geq N}(0) \\ &= \lim_{T \rightarrow \infty} i \int_0^T P^+ e^{-it\Delta} P_{\geq N} F(u(t)) dt - i \lim_{T \rightarrow \infty} \int_{-T}^0 P^- e^{-it\Delta} P_{\geq N} F(u(t)) dt, \end{aligned} \quad (66)$$

where the limit is to be interpreted as a weak limit in L^2 . However, this representation is not useful for $|x|$ small because the kernels of $P^\pm$ have a logarithmic singularity at $x = 0$. To deal with this, we will use a different representation for $|x| \leq N^{-1}$, namely

$$u_{\geq N}(0) = \lim_{T \rightarrow \infty} i \int_0^T e^{-it\Delta} P_{\geq N} F(u(t)) dt, \quad (67)$$

also as a weak limit. To deal with the poor nature of these limits, we use the fact that

$$f_T \rightarrow f \text{ weakly along a subsequence} \Rightarrow \|f\| \leq \limsup_{T \rightarrow \infty} \|f_T\|, \quad (68)$$

or equivalently, that the unit ball is weakly closed.

Despite different representations will be used depending on the size of $|x|$, some estimates can be dealt with in a uniform manner. The first such example is a bound on integrals over short times.

Lemma 6.1 (Local estimate). *For any $\eta > 0$, there exists $\delta = \delta(u, \eta) > 0$ such that*

$$\left\| \int_0^\delta e^{-it\Delta} P_{\geq N} F(u(t)) dt \right\|_{L_x^2} \lesssim_u \eta \mathcal{M}\left(\frac{N}{8}\right),$$

provided that N is sufficiently large depending on u and η . An analogous estimate holds for integration over $[-\delta, 0]$ and after pre-multiplication by $P^\pm$ (they are bounded operators on L_x^2).

Proof. By Strichartz estimates, it suffices to prove

$$\|P_{\geq N} F(u)\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} \lesssim_u \eta \mathcal{M}\left(\frac{N}{8}\right)$$

for any interval J with length $|J| \leq \delta$ and all sufficiently large N depending on u and η .

From (58), there exists $N_0 = N_0(u, \eta)$ such that

$$\|u_{\geq N_0}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} \leq \eta^2. \quad (69)$$

Let $N > 8N_0$. We decompose

$$u = u_{\geq \frac{N}{8}} + u_{N_0 \leq \cdot \leq \frac{N}{8}} + u_{< N_0}.$$

Any term in the resulting expansion of $P_{\geq N} F(u)$ that doesn't contain the factor of $u_{\geq \frac{N}{8}}$ vanishes.

At first, we consider the terms with two factors of the form $u_{< N_0}$. Using Hölder's inequality, Hardy-Littlewood-Sobolev inequality, (57) and Lemma 2.1,

$$\begin{aligned} \|(V * (u_{> \frac{N}{8}} u_{< N_0})) u_{< N_0}\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} &\lesssim \|V * (u_{> \frac{N}{8}} u_{< N_0})\|_{L_t^4 L_x^{\frac{2d}{3}}(J \times \mathbb{R}^d)} \|u_{< N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{> \frac{N}{8}} u_{< N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \|u_{< N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{> \frac{N}{8}}\|_{L_t^\infty L_x^2} \|u_{< N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)}^2 \\ &\lesssim |J|^{\frac{1}{2}} N_0 \mathcal{M}\left(\frac{N}{8}\right) \end{aligned}$$

and

$$\begin{aligned} \|(V * u_{< N_0}^2) u_{> \frac{N}{8}}\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} &\lesssim \|V * u_{< N_0}^2\|_{L_t^2 L_x^d(J \times \mathbb{R}^d)} \|u_{> \frac{N}{8}}\|_{L_t^\infty L_x^2(\mathbb{R} \times \mathbb{R}^d)} \\ &\lesssim \|u_{< N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)}^2 \mathcal{M}\left(\frac{N}{8}\right) \\ &\lesssim |J|^{\frac{1}{2}} N_0 \mathcal{M}\left(\frac{N}{8}\right). \end{aligned}$$

Choosing δ sufficiently small depending on η and N_0 , we see that they are acceptable.

It remains only to consider those components of $P_{\geq N}F(u)$ which involve $u_{\geq \frac{N}{8}}$ and at least one of the other terms is not $u_{< N_0}$.

$$\begin{aligned} \|V * (u_{> \frac{N}{8}} u_{\geq N_0})u\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} &\lesssim \|V * (u_{> \frac{N}{8}} u_{\geq N_0})\|_{L_t^4 L_x^{\frac{2d}{3}}(J \times \mathbb{R}^d)} \|u\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{> \frac{N}{8}}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \|u_{\geq N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \langle |J| \rangle^{\frac{1}{4}}. \end{aligned}$$

By (61), we get

$$\|u_{> N_0}\|_{L_t^2 L_x^{\frac{2d}{d-2}}(J \times \mathbb{R}^d)} \lesssim \langle |J| \rangle^{\frac{1}{2}}.$$

Therefore, by interpolation between this and (69), we have

$$\|u_{> N_0}\|_{L_t^4 L_x^{\frac{2d}{d-1}}(J \times \mathbb{R}^d)} \lesssim \eta \langle |J| \rangle^{\frac{1}{4}}.$$

Thus, we obtain that

$$\|V * (u_{> \frac{N}{8}} u_{\geq N_0})u\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} \lesssim \eta \mathcal{M}\left(\frac{N}{8}\right) \langle |J| \rangle^{\frac{1}{2}}.$$

Similarly, we can estimate

$$\begin{aligned} \|V * (u_{> \frac{N}{8}} u)u_{\geq N_0}\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)} &\lesssim \|V * (u_{> \frac{N}{8}} u)\|_{L_t^2 L_x^d(J \times \mathbb{R}^d)} \|u_{\geq N_0}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{> \frac{N}{8}} u\|_{L_t^2 L_x^{\frac{d}{d-1}}(J \times \mathbb{R}^d)} \|u_{\geq N_0}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \\ &\lesssim \|u_{> \frac{N}{8}}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \|u_{\geq N_0}\|_{L_t^\infty L_x^2(J \times \mathbb{R}^d)} \|u\|_{L_t^2 L_x^{\frac{2d}{d-2}}(J \times \mathbb{R}^d)} \\ &\lesssim \eta^2 \mathcal{M}\left(\frac{N}{8}\right) \langle |J| \rangle^{\frac{1}{2}}, \end{aligned}$$

where the last inequality comes from (69). Another term $\|V * (u_{\geq N_0} u)u_{> \frac{N}{8}}\|_{L_t^2 L_x^{\frac{2d}{d+2}}(J \times \mathbb{R}^d)}$ can be similarly estimated. $\square$

We now turn our attention to $|t| \geq \delta$. In this case we make the decomposition

$$P_{\geq N} = \sum_{M \geq N} P_M \tilde{P}_M,$$

where $\tilde{P}_M := P_{M/2} + P_M + P_{2M}$. In this way, (67) becomes

$$\begin{aligned} u_{\geq N}(0, x) &= i \int_0^\delta e^{-it\Delta} P_{\geq N} F(u(t)) dt \\ &\quad + \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_\delta^T \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt, \end{aligned} \quad (70)$$

which we will use when $|x| \leq N^{-1}$. The analogous reformulation of (66), namely

$$\begin{aligned} u_{\geq N}(0, x) &= i \int_0^\delta P^+ e^{-it\Delta} P_{\geq N} F(u(t)) dt - i \int_{-\delta}^0 P^- e^{-it\Delta} P_{\geq N} F(u(t)) dt \\ &\quad + \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_\delta^T \int_{\mathbb{R}^d} [P_M^+ e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt \\ &\quad - \lim_{T \rightarrow \infty} \sum_{M \geq N} i \int_{-T}^{-\delta} \int_{\mathbb{R}^d} [P_M^- e^{-it\Delta}](x, y) [\tilde{P}_M F(u(t))](y) dy dt, \end{aligned} \quad (71)$$

will be used when $|x| > N^{-1}$.

The next lemma bounds the integrals over the significant region $|y| \gtrsim M|t|$. Let χ_k denote the characteristic function of the set

$$\{(t, y) : 2^k \delta \leq |t| \leq 2^{k+1} \delta, |y| \gtrsim M|t|\}.$$

Lemma 6.2 (Main contribution). *Let $\eta > 0$ be a small number and let δ be as in Lemma 6.1. Then*

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{2^k \delta}^{2^{k+1} \delta} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \lesssim_u \eta \mathcal{M}\left(\frac{N}{64}\right) \quad (72)$$

for all N sufficiently large depending on u and η . An analogous estimate holds with P_M replaced by P_M^+ and P_M^- ; moreover, the time integrals may be taken over $[-2^{k+1} \delta, -2^k \delta]$.

Proof. By Strichartz estimates, we get

$$\begin{aligned} &\left\| \int_{2^k \delta}^{2^{k+1} \delta} \int_{\mathbb{R}^d} [P_M e^{-it\Delta}](x, y) \chi_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \\ &\lesssim \left\| \chi_k(V * |u|^2) \tilde{P}_{> \frac{M}{8}} u \right\|_{L_t^1 L_y^2} + \left\| \chi_k u \tilde{P}_{> \frac{M}{8}} (V * |u|^2) \right\|_{L_t^2 L_y^{\frac{2d}{d+2}}}. \end{aligned}$$

We first consider

$$\begin{aligned} \left\| \chi_k(V * |u|^2) \tilde{P}_{> \frac{M}{8}} u \right\|_{L_t^1 L_y^2} &\lesssim \left\| \chi_k(V * |u|^2) \right\|_{L_t^1 L_y^\infty} \left\| P_{> \frac{M}{64}} u \right\|_{L_t^\infty L_y^2} \\ &\lesssim \left\| \chi_k(y) \int_{|x-y| \geq \frac{|y|}{2}} \frac{|u(x)|^2}{|y-x|^2} dx \right\|_{L_t^1 L_y^\infty} \mathcal{M}\left(\frac{N}{64}\right) \\ &\quad + \left\| \chi_k(y) \int_{|x-y| < \frac{|y|}{2}} \frac{|u(x)|^2}{|y-x|^2} dx \right\|_{L_t^1 L_y^\infty} \mathcal{M}\left(\frac{N}{64}\right). \end{aligned}$$

On one hand,

$$\begin{aligned} \left\| \chi_k(y) \int_{|x-y| \geq \frac{|y|}{2}} \frac{|u(x)|^2}{|y-x|^2} dx \right\|_{L_t^1 L_y^\infty} &\lesssim \left\| \chi_k(y) |y|^{-2} \right\|_{L_t^1 L_y^\infty} \|u\|_{L_t^\infty L_y^2} \\ &\lesssim_u (M 2^k \delta)^{-2} (2^k \delta). \end{aligned}$$

On the other hand, by weighted Strichartz estimates, Hölder's inequality and (62), taking $p = \frac{d}{2} - \frac{1}{4}$, $q = 2d - \frac{3}{2}$ and $\theta = \frac{q}{2p} = \frac{4d-3}{2d-1} \leq 2$, we have

$$\begin{aligned}
& \left\| \chi_k(y) \int_{|x-y| < \frac{|y|}{2}} \frac{|u(x)|^2}{|y-x|^2} dx \right\|_{L_t^1 L_y^\infty} \\
& \lesssim \left\| \chi_k(y) |y|^{-\frac{2(d-1)\theta}{q}} \int_{|x-y| \leq \frac{|y|}{2}} \frac{|y|^{\frac{2(d-1)\theta}{q}} |u(x)|^2}{|x-y|^2} dx \right\|_{L_t^1 L_y^\infty} \\
& \lesssim \left\| \chi_k(y) |y|^{-\frac{2(d-1)\theta}{q}} \int_{|x-y| \leq \frac{|y|}{2}} \frac{|x|^{\frac{2(d-1)\theta}{q}} |u(x)|^2}{|x-y|^2} dx \right\|_{L_t^1 L_y^\infty} \\
& \lesssim \left\| \chi_k(y) |y|^{-\frac{2(d-1)\theta}{q}} \left\| 1_{|\cdot| \leq \frac{|y|}{2}} \frac{1}{|\cdot|^2} \right\|_{L_x^p} \left\| |x|^{\frac{2(d-1)\theta}{q}} |u| \right\|_{L_x^{\frac{2q}{q-4}}}^\theta \|u\|_{L_x^2}^{2-\theta} \right\|_{L_t^1 L_y^\infty} \\
& \lesssim \left\| \chi_k(y) |y|^{-\frac{2(d-1)\theta}{q}} |y|^{\frac{d-2p}{p}} \left\| 1_{|\cdot| \leq \frac{|y|}{2}} \frac{1}{|\cdot|^2} \right\|_{L_t^{\frac{2p}{2p-1}} L_y^\infty} \left\| |x|^{\frac{2(d-1)\theta}{q}} |u| \right\|_{L_t^q L_x^{\frac{2q}{q-4}}}^\theta \|u\|_{L_t^\infty L_x^2}^{2-\theta} \right\|_{L_t^1 L_y^\infty} \\
& \lesssim_u \left(M 2^k \delta \right)^{-\frac{d-1}{p} + \frac{d-2p}{p}} (2^k \delta)^{\frac{2p-1}{2p}} \langle 2^k \delta \rangle^{\frac{\theta}{q}} \lesssim_u M^{-\frac{2p-1}{p}} (2^k \delta)^{\frac{1-p}{p}}.
\end{aligned}$$

Therefore, we have

$$\left\| \chi_k u_{>\frac{M}{8}} (V * |u|^2) \right\|_{L_t^1 L_y^2} \lesssim M^{-\frac{2(2d-3)}{2d-1}} 2^{-\frac{2d-5}{2d-1}k} \delta^{-\frac{2d-5}{2d-1}} \mathcal{M}\left(\frac{N}{64}\right). \quad (73)$$

At last, by means of Bernstein, weighted Strichartz estimates and (62), we have

$$\begin{aligned}
& \left\| \chi_k u_{>\frac{M}{8}} \tilde{P}_{>\frac{M}{8}} (V * |u|^2) \right\|_{L_t^2 L_y^{\frac{2d}{d+2}}} \\
& \lesssim \left\| \chi_k u \right\|_{L_t^2 L_y^\infty} \left\| \tilde{P}_{>\frac{M}{8}} (V * |u|^2) \right\|_{L_t^\infty L_y^{\frac{2d}{d+2}}} \\
& \lesssim M^{-\frac{d-2}{2}} \left\| \chi_k |y|^{-\frac{d-1}{2}} \right\|_{L_t^4 L_y^\infty} \left\| |y|^{\frac{d-1}{2}} u \right\|_{L_t^4 L_y^\infty} \left\| \tilde{P}_{>\frac{M}{8}} (|u|^2) \right\|_{L_t^\infty L_y^1} \\
& \lesssim M^{-\frac{d-2}{2}} (M 2^k \delta)^{-\frac{d-1}{2}} (2^k \delta)^{\frac{1}{4}} \langle 2^k \delta \rangle^{\frac{1}{4}} \|P_{>\frac{M}{64}} u\|_{L_t^\infty L_y^2} \|u\|_{L_t^\infty L_y^2} \\
& \lesssim_u M^{-\frac{d-2}{2}} (M 2^k \delta)^{-\frac{d-1}{2}} (2^k \delta)^{\frac{1}{4}} \langle 2^k \delta \rangle^{\frac{1}{4}} \mathcal{M}\left(\frac{N}{64}\right) \\
& = M^{-\frac{2d-3}{2}} 2^{-\frac{d-2}{2}k} \delta^{-\frac{d-2}{2}} \mathcal{M}\left(\frac{N}{64}\right).
\end{aligned}$$

Thus the LHS of (72) can be bounded by

$$\left(N^{-\frac{2(2d-3)}{2d-1}} \delta^{-\frac{2d-5}{2d-1}} + N^{-\frac{2d-3}{2}} \delta^{-\frac{d-2}{2}} \right) \mathcal{M}\left(\frac{N}{64}\right).$$

This is acceptable as long as we choose N sufficiently large depending on δ and η . $\square$

Now we turn to the integration over the region of (t, y) where $|y| \ll M|t|$. In [12], the bounds of the kernels of the propagators have been shown to be

$$|P_M e^{-it\Delta}(x, y)| + |P_M^\pm e^{-it\Delta}(x, y)| \lesssim \frac{1}{(M^2|t|)^{50d}} K_M(x, y) \quad (74)$$

where

$$K_M(x, y) := \frac{M^d}{\langle M(x - y) \rangle^{50d}} + \frac{M^d}{\langle Mx \rangle^{\frac{d-1}{2}} \langle My \rangle^{\frac{d-1}{2}} \langle M|x| - M|y| \rangle^{50d}}.$$

Furthermore, by Schur's test, it is the kernel of a bounded operator on $L_x^2(\mathbb{R}^d)$.

Let $\tilde{\chi}_k$ be the characteristic function of the set

$$\{(t, y) : 2^k \delta \leq |t| \leq 2^{k+1} \delta, |y| \ll M|t|\}.$$

Lemma 6.3 (The tail). *Let $\eta > 0$ be a small number and let δ be as in Lemma 6.1. Then*

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{2^k \delta}^{2^{k+1} \delta} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2 |t|)^{50d}} \tilde{\chi}_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \lesssim_u \eta \mathcal{M}\left(\frac{N}{8}\right)$$

for all N sufficiently large depending on u and η .

Proof. By Strichartz estimates, Hardy-Littlewood-Sobolev inequality, Hölder inequality and (61), we have

$$\begin{aligned} & \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2 |t|)^{50d}} \tilde{\chi}_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \\ & \lesssim (M^2 2^k \delta)^{-50d} \|\tilde{\chi}_k \tilde{P}_M F(u)\|_{L_t^1 L_x^2} \\ & \lesssim (M^2 2^k \delta)^{-50d} \|u_{\geq \frac{M}{8}}\|_{L_t^\infty L_x^2} \|u\|_{L_t^2 L_x^{\frac{2d}{d-2}}([2^k \delta, 2^{k+1} \delta] \times \mathbb{R}^d)}^2 \\ & \lesssim (M^2 2^k \delta)^{-50d} \mathcal{M}\left(\frac{N}{8}\right) \langle 2^k \delta \rangle. \end{aligned}$$

Summing over $k \geq 0$ and $M \geq N$, we get

$$\sum_{M \geq N} \sum_{k=0}^{\infty} \left\| \int_{\mathbb{R}} \int_{\mathbb{R}^d} \frac{K_M(x, y)}{(M^2 |t|)^{50d}} \tilde{\chi}_k(t, y) [\tilde{P}_M F(u(t))](y) dy dt \right\|_{L_x^2} \lesssim_u (N^2 \delta)^{-49d} \mathcal{M}\left(\frac{N}{8}\right).$$

The claim follows by choosing N sufficiently large depending on δ and η . $\square$

Proof of Proposition 6.1. Naturally, we may bound $\|u_{\geq N}\|_{L^2}$ by separately bounding the L^2 norm on the ball $\{|x| \leq N^{-1}\}$ and on its complement. On the ball we use (70) while outside the ball we use (71). Invoking (68) and the triangle inequality, we reduce the proof to bounding certain integrals. The integrals over short times were estimated in Lemma 6.1. For $|t| \geq \delta$, we further partition the region of integration into two main pieces. The first piece, where $|y| \gtrsim M|t|$, was dealt with in Lemma 6.2. The remaining piece, $|y| \ll M|t|$, can be estimated by combining (74) with Lemma 6.3.

7 Double high-to-low frequency cascade

In this section, we use the additional regularity provided by Theorem 1.5 to preclude double high-to-low frequency cascade solutions. We need the following lemma:

Lemma 7.1 (Gagliardo-Nirenberg inequality of convolution type, [16]). *Let $V(x) = |x|^{-2}$ and*

$$\|u\|_{L^V} := \left(\iint_{\mathbb{R}^d \times \mathbb{R}^d} |u(x)|^2 V(x-y) |u(y)|^2 dx dy \right)^{\frac{1}{4}}, \quad (75)$$

then

$$\|u\|_{L^V}^4 \leq \frac{2}{\|Q\|_{L^2}^2} \|u\|_{L^2}^2 \|\nabla u\|_{L^2}^2. \quad (76)$$

Theorem 7.1 (Absence of double cascades). *There are no non-zero global spherically symmetric solutions to (1) that are double high-to-low frequency cascades in the sense of Theorem 1.3.*

Proof. Suppose there exists such a solution u . By Theorem 1.5, u lies in $C_t^0 H_x^1(\mathbb{R} \times \mathbb{R}^d)$. Hence the energy

$$E(u(t)) := \frac{1}{2} \int_{\mathbb{R}^d} |\nabla u(t, x)|^2 dx + \frac{\mu}{4} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|u(x)|^2 |u(y)|^2}{|x-y|^2} dx dy$$

is finite and conserved. As we have $M(u) < M(Q)$ in the focusing case, Lemma 7.1 gives

$$\|\nabla u(t)\|_{L_x^2(\mathbb{R}^d)}^2 \sim_u E(u) \sim_u 1 \quad (77)$$

for all $t \in \mathbb{R}$. Since

$$\liminf_{t \rightarrow -\infty} N(t) = \liminf_{t \rightarrow +\infty} N(t) = 0,$$

there are two time sequences along which $N(t) \rightarrow 0$.

Let $\eta > 0$ be arbitrary. By Definition 1.4, we can find $C = C(\eta, u) > 0$ such that

$$\int_{|\xi| \geq CN(t)} |\hat{u}(t, \xi)|^2 d\xi \leq \eta^2$$

for all t . Meanwhile, by Theorem 1.5, $u \in C_t^0 H_x^s(\mathbb{R} \times \mathbb{R}^2)$ for some $s > 1$. Thus,

$$\int_{|\xi| \geq CN(t)} |\xi|^{2s} |\hat{u}(t, \xi)|^2 d\xi \lesssim_u 1$$

for all t and some $s > 1$. By Hölder inequality, we obtain

$$\int_{|\xi| \geq CN(t)} |\xi|^2 |\hat{u}(t, \xi)|^2 d\xi \lesssim_u \eta^{2(s-1)/s}.$$

On the other hand, from mass conservation and Plancherel's theorem, we obtain

$$\|\nabla u(t)\|_{L_x^2(\mathbb{R}^d)} \lesssim_u \eta^{(s-1)/s} + CN(t)$$

for all t . As $\eta > 0$ is arbitrary and there exists a sequence of times $t_n \rightarrow \pm\infty$ such that $N(t_n) \rightarrow 0$, we conclude that $\|\nabla u(t_n)\|_{L_x^2(\mathbb{R}^d)} \rightarrow 0$, as $n \rightarrow \infty$. This contradicts with (77). $\square$

8 Death of solitons

Let

$$M_a(t) := 2\Im \int_{\mathbb{R}^d} \bar{u}(t, x) \vec{a}(x) \cdot \nabla u(t, x) dx,$$

then we have (see [14] for similar calculation)

$$\begin{aligned} \partial_t M_a(t) = & - \int_{\mathbb{R}^d} \Delta(\partial_j a_j) |u(t, x)|^2 dx + 4\Re \int_{\mathbb{R}^d} \partial_k a_j u_j u_k dx \\ & - \mu \iint_{\mathbb{R}^d \times \mathbb{R}^d} (\vec{a}(x) - \vec{a}(y)) \cdot \nabla V(x - y) |u(t, x)|^2 |u(t, y)|^2 dx dy. \end{aligned}$$

Lemma 8.1 (Localized virial identity). *Let $\vec{a}(x) = x\psi(\frac{|x|}{R})$, where ψ is a smooth function and $\psi(r) = 1$ when $r \leq 1$; $\psi = 0$ when $r \geq 2$. Then we get*

$$\partial_t M_a(t) = 8E(u(t))$$

$$- \int_{\mathbb{R}^d} \left[\frac{d^2 - 1}{R|x|} \psi'(\frac{|x|}{R}) + \frac{2d + 1}{R^2} \psi''(\frac{|x|}{R}) + \frac{|x|}{R^3} \psi'''(\frac{|x|}{R}) \right] |u(t, x)|^2 dx \quad (78)$$

$$+ 4 \int_{\mathbb{R}^d} \left[\psi(\frac{|x|}{R}) - 1 + \frac{|x|}{R} \psi'(\frac{|x|}{R}) \right] |\nabla u(t, x)|^2 dx \quad (79)$$

$$+ 2\mu \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left[x\psi(\frac{|x|}{R}) - y\psi(\frac{|y|}{R}) - (x - y) \right] \cdot \frac{x - y}{|x - y|^4} |u(t, x)|^2 |u(t, y)|^2 dx dy. \quad (80)$$

Proposition 8.1. *There are no non-zero global spherically symmetric solutions to (1) that are soliton-like in the sense of Theorem 1.3.*

Proof. Assume to the contrary that there is such a solution u . Then by Theorem 1.5, $u \in C_t^0 H_x^s$ for some $s > 1$. In particular,

$$|M_a(t)| \lesssim_u R. \quad (81)$$

Note that in the focusing case, $M(u) < M(Q)$. As a consequence, Lemma 7.1 gives

$$E(u) \gtrsim_u \int_{\mathbb{R}^d} |\nabla u(t, x)|^2 dx > 0. \quad (82)$$

We will show that (78)-(80) constitute only a small fraction of $E(u)$. Combining this fact with Lemma 8.1, we conclude $\partial_t M_a(t) \gtrsim E(u) > 0$, which contradicts with (81). As in [12], (78) and (79) can be bounded by R^{-2} and $\eta^{\frac{s-1}{s}} + \eta$, respectively, where $\eta > 0$ is a small number to be chosen later. The rest of this section is devoted to estimating (80).

By Definition 1.4 and the fact that $N(t) = 1$ for all $t \in \mathbb{R}$, if R is sufficiently large depending on u and η , then

$$\int_{|x| \geq \frac{R}{4}} |u(t, x)|^2 dx \leq \eta \quad (83)$$

for all $t \in \mathbb{R}$. Let χ denote a smooth cutoff to the region $|x| \geq \frac{R}{2}$, chosen so that $\nabla \chi$ is bounded by R^{-1} and supported where $|x| \sim R$. By Lemma 7.1, (82) and (83), we have

$$\begin{aligned}
(80) &\leq C \iint_{\substack{|x| \geq R \\ |y| \geq R}} \left[x(\psi(\frac{|x|}{R}) - 1) - y(\psi(\frac{|y|}{R}) - 1) \right] \cdot \frac{x-y}{|x-y|^4} |u(x)|^2 |u(y)|^2 dx dy \\
&\quad + C \iint_{\substack{|x| \geq R \\ |y| \leq R}} |x(\psi(\frac{|x|}{R}) - 1)| \frac{|u(x)|^2 |u(y)|^2}{|x-y|^3} dx dy \\
&\quad + C \iint_{\substack{|y| \geq R \\ |x| \leq R}} |y(\psi(\frac{|y|}{R}) - 1)| \frac{|u(x)|^2 |u(y)|^2}{|x-y|^3} dx dy \\
&:= I + II + III.
\end{aligned}$$

On one hand,

$$\begin{aligned}
I &\leq C \iint_{\substack{|x| \geq R \\ |y| \geq R}} \frac{|\chi(x)u(x)|^2 |\chi(y)u(y)|^2}{|x-y|^2} dx dy \\
&\lesssim \|\chi u\|_{L_x^2}^2 \|\nabla(\chi u)\|_{L_x^2}^2 \lesssim_u \eta.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
II &\leq C \iint_{\substack{R < |x| \leq 2R \\ |y| \leq R}} \left| x(\psi(\frac{|x|}{R}) - 1) \right| \frac{|\chi(x)u(x)|^2 |u(y)|^2}{|x-y|^3} dx dy \\
&\quad + \iint_{\substack{|x| > 2R \\ |y| \leq R}} |x| \frac{|\chi(x)u(x)|^2 |u(y)|^2}{|x-y|^3} dx dy \\
&\leq C \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|\chi(x)u(x)|^2 |u(y)|^2}{|x-y|^3} dx dy + \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|\chi(x)u(x)|^2 |u(y)|^2}{|x-y|^2} dx dy \\
&\lesssim \|\chi u\|_{L_x^2} \|\nabla(\chi u)\|_{L_x^2} \|\nabla u\|_{L_x^2}^2 + \|\chi u\|_{L_x^2} \|\nabla(\chi u)\|_{L_x^2} \|u\|_{L_x^2} \|\nabla u\|_{L_x^2} \\
&\lesssim_u \eta^{1/2}.
\end{aligned}$$

III can be estimated similarly. Choosing η sufficiently small depending on u and R sufficiently large depending on u and η , we obtain

$$(78) + (79) + (80) \leq \frac{1}{100} E(u).$$

This completes the proof. $\square$

Acknowledgements: The authors thank the referees and the associated editor for their invaluable comments and suggestions which helped improve the paper greatly. C. Miao and G.Xu were partly supported by the NSF of China (No.10725102, No.10726053), and L.Zhao was supported by China postdoctoral science foundation project.

References

- [1] P. Begout and A. Vargas, Mass concentration phenomena for the L^2 -critical nonlinear Schrödinger equation, *Tran. Amer. Math. Soc.* **359** (2007), 5257-5282.
- [2] H. Bahouri and P. Gérard, High frequency approximation of solutions to critical nonlinear wave equations, *Amer. J. Math.* **121** (1999), 131-175.
- [3] J. Bourgain, Global well-posedness of defocusing critical nonlinear Schrödinger equation in the radial case, *J. Amer. Math. Soc.* **12** (1999), 145-171.
- [4] J. Bourgain, Refinements of Strichartz inequality and applications to 2d-NLS with critical nonlinearity, *Int. Math. Res. Not.* (1998), 253-284.
- [5] T. Cazenave and F. B. Weissler, The Cauchy problem for the critical nonlinear Schrödinger equation in H^s , *Nonl. Anal. - TMA* **14** (1990), 807-836.
- [6] J. Colliander, M. Keel, G. Staffilani, H. Takaoka and T. Tao, Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in $\mathbb{R}^3$, *Ann. Math.* **166** (2007), 1-100.
- [7] Yu. B. Gaididei, K. Ø. Rasmussen, P. L. Christiansen, Nonlinear excitations in two-dimensional molecular structures with impurities, *Phys. Rev. E* **52**, 2951-2962.
- [8] C. E. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energy-critical, focusing, non-linear Schrödinger equation in the radial case, *Invent. Math.* **166** (2006), 645-675.
- [9] S. Keraani, On the defect of compactness for the Strichartz estimates of the Schrödinger equations, *J. Diff. Equa.* **175** (2001), 353-392.
- [10] S. Keraani, On the blow up phenomenon of the critical nonlinear Schrödinger equation, *J. Func. Anal.* **235** (2006), 171-192.
- [11] R. Killip, T. Tao and M. Visan, The cubic nonlinear Schrödinger equation in two dimensions with radial data, *arXiv: 0707.3188*.
- [12] R. Killip, M. Visan and X. Zhang, The mass-critical nonlinear Schrödinger equation with radial data in dimensions three and higher, *arXiv:0708.0849*.
- [13] C. Miao, G. Xu and L. Zhao, The Cauchy problem of the Hartree equation, *J. PDE* **21** (2008), 22-44.
- [14] C. Miao, G. Xu and L. Zhao, Global well-posedness and scattering for the energy-critical, defocusing Hartree equation for radial data, *J. Func. Anal.* **253** (2007), 605-627.
- [15] C. Miao, G. Xu and L. Zhao, Global well-posedness and scattering for the energy-critical, defocusing Hartree equation in $\mathbb{R}^{1+n}$, *arXiv:0707.3254*.

- [16] C. Miao, G. Xu and L. Zhao, On the blow up phenomenon for the L^2 -critical focusing Hartree equation in three dimensions, arXiv:0708.2614.
- [17] F. Merle and L. Vega, Compactness at blow-up time for L^2 solutions of the critical nonlinear Schrödinger equation in 2D, Int. Math. Res. Not. **8** (1998), 399-425.
- [18] L. P. Pitaevskii, Vortex lines in an imperfect Bose gas, Sov. Phys. JETP **13**, 451-454.
- [19] S. Shao, Sharp linear and bilinear restriction estimates for the paraboloid in the cylindrically symmetric case, arXiv:0706.3759.
- [20] M. Visan, The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions, Duke. Math. J. **138** (2007), 281-374.
- [21] T. Tao, Nonlinear Dispersive Equations: local and global analysis, CBMS Regional Series in Mathematics, No. 106, 2006
- [22] T. Tao, M. Visan and X. Zhang, Minimal-mass blowup solutions of the mass-critical NLS, arXiv:math/0609690v2.
- [23] T. Tao, M. Visan and X. Zhang, Global well-posedness and scattering for the mass-critical nonlinear Schrödinger equation for radial data in high dimensions, Duke. Math. J. **140** (2007), 165-202.